

CD84 (PN0235) Nb-FC recombinant antibody

CatalogNo: YA0211 **Recombinant** 

Key Features

Reactivity

- Human

Applications

- ELISA,FC

Recommended Dilution Ratios

ELISA 1:5000-100000

Flow Cyt 1-2µg/Test

Storage

Storage* -15°C to -25°C/1 year(Avoid freeze / thaw cycles)**Formulation** Phosphate-buffered solution

Basic Information

Source Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell**Purification** Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell**Clone Number** PN0235


Immunogen Information

Immunogen Purified recombinant Human CD84**Specificity** This recombinant monoclonal antibody can detects endogenous levels of CD84 protein.

Target Information

Gene name	CD84 SLAMF5		
Protein Name	SLAM family member 5 (Cell surface antigen MAX.3) (Hly9-beta) (Leukocyte differentiation antigen CD84) (Signaling lymphocytic activation molecule 5) (CD antigen CD84)		
	Organism	Gene ID	UniProt ID
	Human	8832 ;	Q9UIB8 ;
Cellular Localization	Cell membrane ; Single-pass type I membrane protein .		
Tissue specificity	Predominantly expressed in hematopoietic tissues, such as lymph node, spleen and peripheral leukocytes. Expressed in macrophages, B-cells, monocytes, platelets, thymocytes, T-cells and dendritic cells. Highly expressed in memory T-cells. Expressed in mast cells.		
Function	developmental stage:Expression is slightly increased in naive B-cells after the first division. By contrast, expression on memory B-cells decreased with each successive division.,Domain:ITSM (immunoreceptor tyrosine-based switch motif) motif is a cytoplasmic motif which may bind SH2D1A.,Plays a role as adhesion receptor functioning by homophilic interactions and by clustering. Recruits SH2 domain-containing proteins SH2D1A/SAP. Increases proliferative responses of activated T-cells and SH2D1A/SAP does not seem be required for this process. Homophilic interactions enhance interferon gamma/IFNG secretion in lymphocytes and induce platelet stimulation via a SH2D1A/SAP-dependent pathway. May serve as a marker for hematopoietic progenitor cells.,PTM:N-glycosylated.,PTM:Phosphorylated by tyrosine-protein kinase LCK on tyrosine residues following ligation induced by agonist monoclonal antibody. The association with SH2D1A/SAP is dependent of tyrosines phosphorylation of its cytoplasmic domain Phosphorylated on Tyr-296 and Tyr-316 following platelet aggregation.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,subunit:Homodimer; via its extracellular domain. Forms a head to tail dimer with a CD48 molecule from another cell. Interacts with SH2 domain-containing proteins SH2D1A/SAP and SH2D1B/hEAT-2. Interacts with tyrosine-protein phosphatases PTPN6 and PTPN11 via its phosphorylated cytoplasmic domain, and this interaction is blocked by SH2D1A.,tissue specificity:Predominantly expressed in hematopoietic tissues, such as lymph node, spleen and peripheral leukocytes. Expressed in macrophages, B-cells, monocytes, platelets, thymocytes, T-cells and dendritic cells. Highly expressed in memory T-cells.,		

Validation Data



| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**CD84 (PN0235) Nb-
FC recombinant
antibody**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)