

CD366 (PN0210) Nb-FC recombinant antibody

CatalogNo: YA0292 **Recombinant** 

Key Features

Reactivity

- Human

Applications

- ELISA

Recommended Dilution Ratios

ELISA 1:5000-100000

Storage

Storage* -15°C to -25°C/1 year(Avoid freeze / thaw cycles)**Formulation** Phosphate-buffered solution

Basic Information

Source Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell**Purification** Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell**Clone Number** PN0210


Immunogen Information

Immunogen Purified recombinant Human TIM-3**Specificity** This recombinant monoclonal antibody can detects endogenous levels of CD366/TIM-3 protein.

Target Information

Gene name	HAVCR2 TIM3 TIMD3		
Protein Name	Hepatitis A virus cellular receptor 2 (HAVcr-2) (T-cell immunoglobulin and mucin domain-containing protein 3) (TIMD-3) (T-cell immunoglobulin mucin receptor 3) (TIM-3) (T-cell membrane protein 3) (CD antigen CD366)		
	Organism	Gene ID	UniProt ID
	Human	84868 ;	Q8TDQ0 ;
Cellular Localization	Membrane ; Single-pass type I membrane protein . Cell junction . Cell membrane . Localizes to the immunological synapse between CD8+ T-cells and target cells. .		
Tissue specificity	Expressed in T-helper type 1 (Th1) lymphocytes. Expressed on regulatory T (Treg) cells after TCR stimulation. Expressed in dendritic cells and natural killer (NK) cells. Expressed in epithelial tissues. Expression is increased on CD4+ and CD8+ T-cells in chronic hepatitis C virus (HCV) infection. In progressive HIV-1 infection, expression is up-regulated on HIV-1-specific CD8 T-cells.		
Function	Disease:Genetic variation in HAVCR2 may influence susceptibility to rheumatoid arthritis (RA) [MIM:180300]. RA is an inflammatory disease, primarily of the joints, with autoimmune features and a complex genetic component. It is characterized by inflammation of synovial tissues and the formation of rheumatoid pannus, which is capable of eroding adjacent cartilage and bone, causing subsequent joint destruction.,Regulates macrophage activation. Inhibits T-helper type 1 lymphocyte (Th1)-mediated auto- and alloimmune responses and promotes immunological tolerance. May be also involved in T-cell homing. Receptor for LGALS9.,similarity:Belongs to the immunoglobulin superfamily. TIM family.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,tissue specificity:T-helper type 1 lymphocyte (Th1)-specific.,		

Validation Data



Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**CD366 (PN0210) Nb-
FC recombinant
antibody**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)