

## CD16 (PN0225) Nb-FC recombinant antibody

CatalogNo: YA0426 **Recombinant** 

### Key Features

#### Reactivity

- Human

#### Applications

- ELISA,FC

### Recommended Dilution Ratios

**ELISA 1:5000-100000**

**Flow Cyt 1-2µg/Test**

### Storage

**Storage\*** -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

**Formulation** Phosphate-buffered solution

### Basic Information

**Source** Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell

**Purification** Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell

**Clone Number** PN0225

### Immunogen Information


**Immunogen** Purified recombinant Human CD16

**Specificity** This recombinant monoclonal antibody can detects endogenous levels of CD16 protein.

## Target Information

Gene name	FCGR3A CD16A FCG3 FCGR3 IGFR3		
Protein Name	Low affinity immunoglobulin gamma Fc region receptor III-A (IgG Fc receptor III-A) (CD16-II) (CD16a antigen) (Fc-gamma RIII-alpha) (Fc-gamma RIII) (Fc-gamma RIIIa) (FcRIII) (FcRIIIa) (FcgammaRIIIA) (FcR-10) (IgG Fc receptor III-2) (CD antigen CD16a)		
	Organism	Gene ID	UniProt ID
	Human	<a href="#">2214</a> ;	<a href="#">P08637</a> ;
Cellular Localization	Cell membrane ; Single-pass type I membrane protein . Secreted . Exists also as a soluble receptor. .		
Tissue specificity	Expressed in natural killer cells (at protein level) (PubMed:2526846). Expressed in a subset of circulating monocytes (at protein level) (PubMed:2767158).		
Function	Receptor for the Fc region of IgG. Binds complexed or aggregated IgG and also monomeric IgG. Mediates antibody-dependent cellular cytotoxicity (ADCC) and other antibody-dependent responses, such as phagocytosis.,miscellaneous:Encoded by one of two nearly identical genes: FCGR3A (Shown here) and FCGR3B which are expressed in a tissue-specific manner. The Phe-203 in III-A determines the transmembrane domains whereas the Ser-203 in III-B determines the GPI-anchoring.,online information:FCGR3A mutation db,polymorphism:Isoform Val-157 shows a higher binding capacity of IgG1, IgG3 and IgG4 compared with isoform Phe-157. Alleles Leu-66 and Phe-157, and alleles His-66 / Arg-66 and Val-157 are in linkage disequilibrium.,PTM:Glycosylated. Contains high mannose- and complex-type oligosaccharides.,PTM:The soluble form is produced by a proteolytic cleavage.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subcellular location:Exists also as a soluble receptor.,subunit:Exists as a hetero-oligomeric receptor complex with Fc epsilon receptor I gamma subunit and / or the CD3 zeta subunit. Interacts with INPP5D/SHIP1.,tissue specificity:Expressed on natural killer cells, macrophages, subpopulation of T-cells, immature thymocytes and placental trophoblasts.,		

## Validation Data



## Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**CD16 (PN0225) Nb-  
FC recombinant  
antibody**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)