

www.immunoway.com.cn

CD141 (PN0329) Nb-FC recombinant antibody

CatalogNo: YA0558 Recombinant R

Key Features

Reactivity

Human

Applications
• ELISA,FC

Recommended Dilution Ratios

ELISA 1:5000-100000 Flow Cyt 1-2µg/Test

Storage

| Storage* | -15°C to -25°C/1 year(Avoid freeze / thaw cycles) |
|-------------|---|
| Formulation | Phosphate-buffered solution |

Basic Information

| Source | Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell |
|--------------|---|
| Purification | Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell |
| Clone Number | PN0329 |


Immunogen Information

| Immunogen | Purified recombinant Human CD141 |
|-------------|--|
| Specificity | This recombinant monoclonal antibody can detects endogenous levels of CD141 protein. |

Target Information

| Gene name | THBD THRM | | | |
|--------------------------|--|--------------|----------------|--|
| Protein Name | Thrombomodulin (TM) (Fetomodulin) (CD antigen CD141) Organism Gene ID UniProt ID | | | |
| | Human | <u>7056;</u> | <u>P07204;</u> | |
| Cellular Localization | Membrane; Single-pass type I membrane protein. | | | |
| Tissue specificity | Endothelial cells are unique in synthesizing thrombomodulin. | | | |
| Function | Disease:Defects in THBD are the cause of thrombophilia due to thrombomodulin defect (THR-THBDD) [MIM:188040]. THR-THBDD is a hemostatic disorder characterized by a tendency to thrombosis.,Thrombomodulin is a specific endothelial cell receptor that forms a 1:1 stoichiometric complex with thrombin. This complex is responsible for the conversion of protein C to the activated protein C (protein Ca). Once evolved, protein Ca scissions the activated cofactors of the coagulation mechanism, factor Va and factor VIIIa, and thereby reduces the amount of thrombin generated.,online information:Thrombomodulin,online information:Thrombomodulin entry,PTM:N-glycosylated.,PTM:The iron and 2-oxoglutarate dependent 3-hydroxylation of aspartate and asparagine is (R) stereospecific within EGF domains.,similarity:Contains 1 C-type lectin domain.,similarity:Contains 6 EGF-like domains.,tissue specificity:Endothelial cells are unique in synthesizing thrombomodulin., | | | |

Validation Data



Contact information

| Orders: | order.cn@immunoway.com |
|------------|--|
| Support: | support.cn@immunoway.com |
| Telephone: | 400-8787-807(China) |
| Website: | http://www.immunoway.com.cn |
| Address: | 2200 Ringwood Ave San Jose, CA 95131 USA |
| | |



Please scan the QR code to access additional product information: CD141 (PN0329) Nb-FC recombinant antibody For Research Use Only. Not for Use in Diagnostic Procedures.

Immunoway - 3 / 3