

#### www.immunoway.com.cn

# CD38 (PN0518) Nb-FC recombinant antibody

CatalogNo: YA0617 Recombinant R

### Key Features

Reactivity

Human

Applications

ELISA

#### **Recommended Dilution Ratios**

ELISA 1:5000-100000

#### **Storage**

Storage\* -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

Formulation Phosphate-buffered solution

#### **Basic Information**

Source	Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell
Purification	Camel, chimeric fusion of Nanobody (VHH) and mouse $\mbox{IgG1}$ Fc domain , recombinantly produced from 293F cell
Clone Number	PN0518


#### Immunogen Information

Immunogen	Purified recombinant Human CD38
Specificity	This recombinant monoclonal antibody can detects endogenous levels of CD38 protein.

#### **Target Information**

Gene name	CD38		
Protein Name	ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1 (2'-phospho-ADP-ribosyl cyclase) (2'- phospho-ADP-ribosyl cyclase/2'-phospho-cyclic-ADP-ribose transferase) (2'-phospho-cyclic ADP-ribose transferase) (ADP-ribosyl cyclase 1) (ADPRC 1) (Cyclic ADP-ribose hydrolase 1) (cADPr hydrolase 1) (T10) (CD antigen CD38)		
	Organism	Gene ID	UniProt ID
	Human	<u>952;</u>	<u>P28907;</u>
Cellular Localization	Membrane; Single-pass type II membrane protein.		
Tissue specificity	Expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma.		
Function	Catalytic activity:NAD(+) + H(2)O = ADP-ribose + nicotinamide.,developmental stage:Preferentially expressed at both early and late stages of the B and T-cell maturation. It is also detected on erythroid and myeloid progenitors in bone marrow, where the level of surface expression was shown to decrease during differentiation of blast-forming unit E to colony-forming unit E.,enzyme regulation:ATP inhibits the hydrolyzing activity.,Synthesizes cyclic ADP-ribose, a second messenger for glucose-induced insulin secretion. Also has cADPr hydrolase activity. Also moonlights as a receptor in cells of the immune system.,online information:CD38 entry,similarity:Belongs to the ADP-ribosyl cyclase family.,tissue specificity:Expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma.,		

## Validation Data



# **Contact** information

Orders:	order.cn@immunoway.com
Support:	support.cn@immunoway.com
Telephone:	400-8787-807(China)
Website:	http://www.immunoway.com.cn
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: CD38 (PN0518) Nb-FC recombinant antibody For Research Use Only. Not for Use in Diagnostic Procedures.

Immunoway - 3 / 3