

# CD31 (PN0092) Nb-FC recombinant antibody

CatalogNo: YA0620 Recombinant R

### **Key Features**

Reactivity Applications
• Human • FC,ELISA

# **Recommended Dilution Ratios**

ELISA 1:5000-100000 Flow Cyt 1-2µg/Test

### Storage

**Storage\*** -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

**Formulation** Phosphate-buffered solution

## **Basic Information**

**Source** Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain, recombinantly

produced from 293F cell

**Purification** Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain, recombinantly

produced from 293F cell

Clone Number PN0092

### Immunogen Information

Immunogen	Purified recombinant Human CD31
Specificity	This recombinant monoclonal antibody can detects endogenous levels of CD31 protein.

### **Target Information**

#### Gene name

PECAM1

#### **Protein Name**

Platelet endothelial cell adhesion molecule (PECAM-1) (EndoCAM) (GPIIA') (PECA1) (CD antigen CD31)

Organism	Gene ID	UniProt ID
Human	<u>5175;</u>	<u>P16284;</u>

### Cellular Localization


Cell membrane; Single-pass type I membrane protein. Cell surface expression on neutrophils is down-regulated upon fMLP or CXCL8/IL8-mediated stimulation. .; [Isoform Long]: Cell membrane; Single-pass type I membrane protein. Membrane raft. Cell junction . Localizes to the lateral border recycling compartment (LBRC) and recycles from the LBRC to the junction in resting endothelial cells. .; [Isoform Delta15]: Cell junction . Localizes to the lateral border recycling compartment (LBRC) and recycles from the LBRC to the junction in resting endothelial cells.

**Tissue specificity** Expressed on platelets and leukocytes and is primarily concentrated at the borders between endothelial cells (PubMed:18388311, PubMed:21464369). Expressed in human umbilical vein endothelial cells (HUVECs) (at protein level) (PubMed:19342684, PubMed:175838). Expressed on neutrophils (at protein level) (PubMed:175838). Isoform Long predominates in all tissues examined (PubMed:12433657). Isoform Delta12 is detected only in trachea (PubMed:12433657). Isoform Delta14-15 is only detected in lung (PubMed:12433657). Isoform Delta14 is detected in all tissues examined with the strongest expression in heart (PubMed:12433657). Isoform Delta15 is expressed in brain, testis, ovary, cell surface of platelets, human umbilical vein endothelial cells (HUVECs), Jurkat T-cell leukemia, human erythroleukemia (HEL) and U-937 histiocytic lymphoma cell lines (at protein level) (PubMed:12433657, PubMed:18388311).

#### **Function**

This protein is a cell adhesion molecule expressed on platelets and at endothelial cell intercellular junctions., online information: CD31 entry, online information: PECAM-1, online information: The Singapore human mutation and polymorphism database,PTM:Phosphorylated on Ser and Tyr residues after cellular activation., similarity: Contains 6 Ig-like C2-type (immunoglobulin-like) domains., tissue specificity:Long isoform predominates all tissues examined, isoform Delta12 was detected only in trachea and isoform Delta14-15 only in lung, isoform Delta14 was detected in all tissues examined with the strongest expression in heart.,

### Validation Data



### | Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

CD31 (PN0092) Nb-FC recombinant antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents