

#### www.immunoway.com.cn

# Cleaved KLK11 (Ile54) Rabbit pAb

CatalogNo: YC0055 Orthogonal Validated 💽

# Key Features

Host Species • Rabbit	Reactivity <ul> <li>Human,Rat,Mouse,</li> </ul>	<ul><li>Applications</li><li>WB,ELISA</li></ul>
MW • 25kD (Observed)	Isotype • IgG	

#### **Recommended Dilution Ratios**

WB 1:500-1:2000 ELISA 1:10000 Not yet tested in other applications.

### **Storage**

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)		
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		

### **Basic Information**

Clonality Polyclonal

#### Immunogen Information

Immunogen	The antiserum was produced against synthesized peptide derived from human Kallikrein-11. AA range:35-84
Specificity	Cleaved-KLK11 (I54) Polyclonal Antibody detects endogenous levels of fragment of activated KLK11 protein resulting from cleavage adjacent to I54.

Gene name	KLK11				
Protein Name	Kallikrein-11				
	Organism	Gene ID	UniProt ID		
	Human	<u>11012;</u>	<u>Q9UBX7;</u>		
	Mouse		<u>Q9QYN3;</u>		
Cellular Localization	[Isoform 1]: Secreted.; [Isoform 2]: Golgi apparatus.				
Tissue specificity	Expressed in brain, skin and prostate. Isoform 1 is expressed preferentially in brain. Isoform 2 is expressed in prostate. Present in seminal plasma at concentrations ranging from 2 to 37 microg/mL (at protein level).				
Function	Function:Possible multifunctional protease. Efficiently cleaves 'bz-Phe-Arg-4- methylcoumaryl-7-amide', a kallikrein substrate, and weakly cleaves other substrates for kallikrein and trypsin. Cleaves synthetic peptides after arginine but not lysine residues.,PTM:About 40% of KLK11 is inactivated by internal cleavage after Arg-188. This proteolytic inactivation may be effected by plasminogen.,similarity:Belongs to the peptidase S1 family.,similarity:Belongs to the peptidase S1 family. Kallikrein subfamily.,similarity:Contains 1 peptidase S1 domain.,tissue specificity:Expressed in brain, skin and prostate. Isoform 1 is expressed preferentially in brain; isoform 2 in prostate. Present in seminal plasma at concentrations ranging from 2 to 37 microg/mL (at protein level).,				





Western Blot analysis of various cells using Cleaved-KLK11 (I54) Polyclonal Antibody



Western blot analysis of lysates from HeLa cells, treated with etoposide 25uM 24H, using Kallikrein-11 (Cleaved-IIe54) Antibody. The lane on the right is blocked with the synthesized peptide.

## **Contact information**

Orders:order.cn@immunoway.comSupport:support.cn@immunoway.comTelephone:400-8787-807(China)Website:http://www.immunoway.com.cnAddress:2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: Cleaved KLK11 (Ile54) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents