

www.immunoway.com.cn

STAT3 protein

CatalogNo: YD0103

Key Features

Reactivity

Human

Applications
• WB,SDS-PAGE

Recommended Dilution Ratios

Storage

Storage*	-20°C/6 month,-80°C for long storage
Formulation	Liquid in PBS

Basic Information

Purity SDS-PAGE >90%

Immunogen Information

Squence Amino acid: 483-770, with his-MBP tag.

Target Information

Gene name STAT3 APRF

Protein Name STAT3 protein

Organism	Gene ID	UniProt ID
Human	<u>6774;</u>	<u>P40763;</u>

Cellular Localization	Cytoplasm . Nucleus . Shuttles between the nucleus and the cytoplasm. Translocated into the nucleus upon tyrosine phosphorylation and dimerization, in response to signaling by activated FGFR1, FGFR2, FGFR3 or FGFR4. Constitutive nuclear presence is independent of tyrosine phosphorylation. Predominantly present in the cytoplasm without stimuli. Upon leukemia inhibitory factor (LIF) stimulation, accumulates in the nucleus. The complex composed of BART and ARL2 plays an important role in the nuclear translocation and retention of STAT3. Identified in a complex with LYN and PAG1.
Tissue specificity	Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Expressed in naive CD4(+) T cells as well as T-helper Th17, Th1 and Th2 cells (PubMed:31899195).
Function	negative regulation of transcription from RNA polymerase II promoter, eye development, temperature homeostasis,eye photoreceptor cell differentiation, acute inflammatory response, transcription, regulation of transcription, DNA- dependent, regulation of transcription from RNA polymerase II promoter, cell motion, defense response, acute-phase response, inflammatory response, cell surface receptor linked signal transduction, enzyme linked receptor protein signaling pathway, transmembrane receptor protein tyrosine kinase signaling pathway, intracellular signaling cascade, protein kinase cascade, JAK-STAT cascade, sensory organ development, behavior, feeding behavior, response to wounding, response to endogenous stimulus, response to hormone stimulus, negative regulation of biosynthetic process, positive regulation of biosynthetic process, negative regulation of macromolecule biosynthetic process, positive regulation of macromolecule metabolic process, negative regulation of macromolecule metabolic process, negative regulation of gene expression, negative regulation of gene expression, response to organic cyclic substance, negative regulation of transcription, cytokine-mediated signaling pathway, sexual reproduction, neuron differentiation, negative regulation of cellular biosynthetic process, positive regulation of gene expression, response to estradiol stimulus, cellular response to hormone stimulus, carobnydrate homeostasis, response to cytokine stimulus, regulation of growth, regulation of multicellular organism growth, response to drug, homeostatic process, glucose homeostasie, eating behavior, response to eptide hormone stimulus, response to setrogen stimulus, regulation of transcription, positive regulation of transcription, DNA- dependent, positive regulation of transcription, DNA- dependent, positive regulation of transcription, DNA- dependent, positive regulation of transcription from RNA polymerase II promoter, photoreceptor cell differentiation, response to steroid hormone stimulus, eye morphogenesis, m

Validation Data



Contact information

Orders:	order.cn@immunoway.com
Support:	support.cn@immunoway.com
Telephone:	400-8787-807(China)
Website:	http://www.immunoway.com.cn
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **STAT3 protein**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents