

Recombinant Human IL1B protein ,C- His Tag

CatalogNo: YD2203

| Key Features

Reactivity

Human

Recommended Dilution Ratios

Reconstitute in sterile water for a stock solution

Storage

Storage* Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 2 to 8 °C

for one week. Store at -20 to -80 °C for twelve months from the date of receipt.

Formulation Supplied as solution form in PBS pH7.5./ Lyophilized from PBS pH7.5.

I Basic Information

Source E.coli

Purification E.coli

Purity >90% as determined by SDS-PAGE

Immunogen Information

Squence A DNA sequence encoding the human IL1B(Ala117~Ser269) was fused with the C-

terminal His Tag

| Target Information

Protein Name	Organism	Gene ID	UniProt ID	
	Human	<u>3553</u> ;	<u>P01584;</u>	

Cellular Localization

Cytoplasm, cytosol . Secreted . Lysosome . Secreted, extracellular exosome . The precursor is cytosolic (PubMed:15192144). In response to inflammasome-activating signals, such as ATP for NLRP3 inflammasome or bacterial flagellin for NLRC4 inflammasome, cleaved and secreted (PubMed:24201029, PubMed:33377178, PubMed:33883744). Mature form is secreted and released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed:33883744). In contrast, the precursor form is not released, due to the presence of an acidic region that is proteolytically removed by CASP1 during maturation (PubMed:33883744). The secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10 (PubMed:32272059).

Tissue specificity Expressed in activated monocytes/macrophages (at protein level).

Validation Data

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

Recombinant Human IL1B protein ,C- His Tag

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents