

CD319 (CRACC) recombinant protein

CatalogNo: YD3008

| Key Features

Reactivity

- Human

| Storage

Storage* -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

Formulation Phosphate-buffered solution

| Recommended Dilution Ratios

| Basic Information

Source Mammalian cells

Purification Mammalian cells

Purity >90% as determined by SDS-PAGE

| Immunogen Information

Sequence Amino acid:23-226,with human FC tag.

| Target Information

Gene name SLAMF7 CS1 UNQ576/PRO1138

Protein Name	SLAM family member 7		
	Organism	Gene ID	UniProt ID
	Human	57823 ;	Q9NQ25 ;
Cellular Localization	Membrane; Single-pass type I membrane protein.		
Tissue specificity	Expressed in spleen, lymph node, peripheral blood leukocytes, bone marrow, small intestine, stomach, appendix, lung and trachea. Expression was detected in NK cells, activated B-cells, NK-cell line but not in promyelocytic, B-, or T-cell lines. Expressed in monocytes. Isoform 3 is expressed at much lower level than isoform 1.		
Function	Self-ligand receptor of the signaling lymphocytic activation molecule (SLAM) family. SLAM receptors triggered by homo- or heterotypic cell-cell interactions are modulating the activation and differentiation of a wide variety of immune cells and thus are involved in the regulation and interconnection of both innate and adaptive immune response. Activities are controlled by presence or absence of small cytoplasmic adapter proteins, SH2D1A/SAP and/or SH2D1B/EAT-2. Isoform 1 mediates NK cell activation through a SH2D1A-independent extracellular signal-regulated ERK-mediated pathway (PubMed:11698418). Positively regulates NK cell functions by a mechanism dependent on phosphorylated SH2D1B. Downstream signaling implicates PLCG1, PLCG2 and PI3K (PubMed:16339536). In addition to heterotypic NK cells-target cells interactions also homotypic interactions between NK cells may contribute to activation. However, in the absence of SH2D1B, inhibits NK cell function. Acts also inhibitory in T-cells (By similarity). May play a role in lymphocyte adhesion (PubMed:11802771). In LPS-activated monocytes negatively regulates production of pro-inflammatory cytokines (PubMed:23695528).; Isoform 3 does not mediate any NK cell activation.		

Validation Data

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
 to access additional
 product information:
CD319 (CRACC)
recombinant protein