

Purified Mouse Anti Human CD55 Antibody (PTF0063)

CatalogNo: YF0062 **Recombinant** 

Key Features

Host Species

- Mouse

Reactivity

- Human

Applications

- IF,FC

Isotype

- Mouse IgG2a/Kappa

Storage

Storage*

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

Formulation

PBS, pH 7.4

Recommended Dilution Ratios

IF 1:200-1:1000

FC:1 µg per million cells in 100 µl volume. For optimal results, the reagent should be titrated according to each specific application

Basic Information

Clonality

Monoclonal

Clone Number

PTF0063

Immunogen Information

Immunogen

The specific immunogen used to produce this antibody is proprietary information.

Specificity

Endogenous

| Target Information

Gene name CD55

Protein Name Complement decay-accelerating factor


Organism	Gene ID	UniProt ID
Human	1604 ;	P08174 ;

Cellular Localization [Isoform 1]: Cell membrane; Single-pass type I membrane protein.; [Isoform 2]: Cell membrane; Lipid-anchor , GPI-anchor.; [Isoform 3]: Secreted .; [Isoform 4]: Secreted .; [Isoform 5]: Secreted .; [Isoform 6]: Cell membrane ; Lipid-anchor , GPI-anchor .; [Isoform 7]: Cell membrane ; Lipid-anchor , GPI-anchor .

Tissue specificity Expressed on the plasma membranes of all cell types that are in intimate contact with plasma complement proteins. It is also found on the surfaces of epithelial cells lining extracellular compartments , and variants of the molecule are present in body fluids and in extracellular matrix.

Function Domain:The first Sushi domain (SCR1) is not necessary for function. SCR2 and SCR4 provide the proper conformation for the active site on SCR3. ,Function:This protein recognizes C4b and C3b fragments that condense with cell-surface hydroxyl or amino groups when nascent C4b and C3b are locally generated during C4 and c3 activation. Interaction of daf with cell-associated C4b and C3b polypeptides interferes with their ability to catalyze the conversion of C2 and factor B to enzymatically active C2a and Bb and thereby prevents the formation of C4b2a and C3bBb , the amplification convertases of the complement cascade. ,online information:Blood group antigen gene mutation database ,online information:CD55 mutation db ,online information:Decay-accelerating factor entry ,online information:Icosahedral capsid structure ,polymorphism:Responsible for the Cromer blood group system. It consists of at least 8 high-incidence (Cr (a) , Tc (a) , Dr (a) , Es (a) , WES (b) , UMC , IFC and GUTI) and three low-incidence (Tc (b) , Tc (c) and WES (a)) antigens that reside on DAF. In the Cromer phenotypes Dr (a-) and Inab there is reduced or absent expression of DAF , respectively. In the case of the Dr (a-) phenotype , a single nucleotide substitution within exon 5 accounts for two changes: a simple amino acid substitution , Leu-199 that is the basis of the antigenic variation , and an alternative splicing event that underlies the decreased expression of DAF in this phenotype. The Inab phenotype is a very rare one in which the red blood cells lack all Cromer system antigens. The red blood cells of individuals with Inab phenotype have a deficiency of DAF , but these individuals are not known to have any associated hematologic or other abnormalities. ,PTM:The Ser/Thr-rich domain is heavily O-glycosylated. ,similarity:Belongs to the receptors of complement activation (RCA) family. ,similarity:Contains 4 Sushi (CCP/SCR) domains. ,subunit:Monomer (major form) and non-disulfide-linked , covalent homodimer (minor form) . Binds to coxsackievirus A21 , coxsackieviruses B1 , B3 and B5 , human enterovirus 70 , human echoviruses 6 , 7 , 11 , 12 , 20 and 21 capsid proteins and acts as a receptor for these viruses. ,tissue specificity:Expressed on the plasma membranes of all cell types that are in intimate contact with plasma complement proteins. It is also found on the surfaces of epithelial cells lining extracellular compartments , and variants of the molecule are present in body fluids and in extracellular matrix. ,

| Validation Data



Human peripheral blood lymphocytes stained with Mouse IgG2a,κ isotype control or purified CD55 (PTF0063), followed by anti-mouse IgGs-PE

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Purified Mouse Anti Human CD55 Antibody (PTF0063)

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)