

Acetyl eIF5A/eIF5A2 (Lys47) Rabbit pAb

CatalogNo: YK0021

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IF, ELISA

MW

- 17kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IF 1:200-1:1000

ELISA 1:20000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized Acetyl-peptide derived from human eIF5A around the Acetylation site of Lys47. AA range:11-60

Specificity

Acetyl-eIF5A/eIF5A2 (K47) Polyclonal Antibody detects endogenous levels of eIF5A/eIF5A2 protein only when acetylated at K47. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):TSkTG

Target Information

Gene name EIF5A2

Protein Name Eukaryotic translation initiation factor 5A-2

Organism	Gene ID	UniProt ID
Human	56648 ;	Q9GZV4 ;
Mouse	208691 ;	Q8BGY2 ;

Cellular Localization


Cytoplasm . Nucleus . Endoplasmic reticulum membrane ; Peripheral membrane protein ; Cytoplasmic side . Nucleus , nuclear pore complex . Hypusine modification promotes the nuclear export and cytoplasmic localization and there was a dynamic shift in the localization from predominantly cytoplasmic to primarily nuclear under apoptotic inducing conditions. .

Tissue specificity Expressed in ovarian and colorectal cancer cell lines (at protein level) . Highly expressed in testis. Overexpressed in some cancer cells.


Function

Function: The precise role of eIF-5A in protein biosynthesis is not known but it functions by promoting the formation of the first peptide bond. , PTM: eIF-5A seems to be the only eukaryotic protein to have an hypusine residue which is a post-translational modification of a lysine by the addition of a butylamino group (from spermidine) . , similarity: Belongs to the eIF-5A family. , tissue specificity: Expressed in ovarian and colorectal cancer cell lines (at protein level) . Highly expressed in testis. Overexpressed in some cancer cells. ,


Validation Data



Immunofluorescence analysis of MCF7 cell. 1, primary Antibody was diluted at 1:100 (4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - AFluor 594 Secondary antibody (catalog No: RS3611) was diluted at 1:500 (room temperature, 50min).



Western blot analysis of L929 lysis using antibody diluted at 1:1000. Secondary antibody (catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from L929 cells, using eIF5A (Acetyl-Lys47) Antibody.

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**Acetyl eIF5A/eIF5A2
(Lys47) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)