

## ATF-5 (Acetyl Lys29) Rabbit pAb

CatalogNo: YK0024

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, ELISA

#### MW

- 30kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:500-1:2000**

**ELISA 1:20000**

**Not yet tested in other applications.**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized Acetyl-peptide derived from human ATF5 around the Acetylation site of Lys29. AA range:1-50

**Specificity** Acetyl-ATF-5 (K29) Polyclonal Antibody detects endogenous levels of ATF-5 protein only when acetylated at K29. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): YGkLP

## Target Information

**Gene name** ATF5

**Protein Name** Cyclic AMP-dependent transcription factor ATF-5

Organism	Gene ID	UniProt ID
Human	<a href="#">22809;</a>	<a href="#">Q9Y2D1;</a>
Mouse	<a href="#">107503;</a>	<a href="#">O70191;</a>
Rat	<a href="#">282840;</a>	<a href="#">Q6P788;</a>

**Cellular Localization**


Cytoplasm . Nucleus . Cytoplasm , cytoskeleton , microtubule organizing center , centrosome . Actively transported to the centrosome and accumulated in the pericentriolar material (PCM) during G1 to M phase via a microtubule-dependent mechanism. During late telophase and cytokinesis , translocates from the centrosome to the midbody. .

**Tissue specificity** Widely expressed with higher expression levels in liver.

**Function**

Function:Transcriptional activator which binds the cAMP response element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3') , a sequence present in many viral and cellular promoters and blocks the differentiation of neuroprogenitor cells into neurons. Its transcriptional activity is enhanced by CCND3 and slightly inhibited by CDK4. ,similarity:Belongs to the bZIP family. ,similarity:Contains 1 bZIP domain. ,subunit:Binds DNA as a dimer. Interacts with PTP4A1/PRL-1 (By similarity) . Interacts with CCND3 , but not with CCND1 or CCND2. ,

## Validation Data



Western Blot analysis of A549 cells using Acetyl-ATF-5 (K29) Polyclonal Antibody. Secondary antibody (catalog#:RS0002) was diluted at 1:20000

## Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: http://www.immunoway.com.cn  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**ATF-5 (Acetyl  
Lys29) Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)