

Acetyl-EPAS-1 (Lys385) Rabbit pAb

CatalogNo: YK0073

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA

MW

- 120kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

ELISA 1:10000-20000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized acetyl-peptide of Acetyl-EPAS-1 (K385)

Specificity Acetyl-EPAS-1 (K385) Polyclonal Antibody detects endogenous levels of Acetyl-EPAS-1 (K385). The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): EkSNF

| Target Information

Gene name EPAS1 BHLHE73 HIF2A MOP2 PASD2

Protein Name Endothelial PAS domain-containing protein 1

Organism	Gene ID	UniProt ID
Human	2034 ;	Q99814 ;
Mouse	13819 ;	P97481 ;
Rat	29452 ;	Q9JHS1 ;

Cellular Localization Nucleus . Nucleus speckle . Colocalizes with HIF3A in the nucleus and speckles. .

Tissue specificity Expressed in most tissues , with highest levels in placenta , lung and heart. Selectively expressed in endothelial cells.

Function Disease:Defects in EPAS1 are the cause of erythrocytosis familial type 4 (ECYT4) [MIM:611783]. ECYT4 is an autosomal dominant disorder characterized by increased serum red blood cell mass , elevated hemoglobin concentration and hematocrit , and normal platelet and leukocyte counts. ,Function:Transcription factor involved in the induction of oxygen regulated genes. Binds to core DNA sequence 5'-[AG]CGTG-3' within the hypoxia response element (HRE) of target gene promoters. Regulates the vascular endothelial growth factor (VEGF) expression and seems to be implicated in the development of blood vessels and the tubular system of lung. May also play a role in the formation of the endothelium that gives rise to the blood brain barrier. Potent activator of the Tie-2 tyrosine kinase expression. Activation seems to require recruitment of transcriptional coactivators such as CREBBP and probably EP300. Interaction with redox regulatory protein APEX seems to activate CTAD. ,PTM:In normoxia , is hydroxylated on Asn-847 by HIF1AN thus probably abrogating interaction with CREBBP and EP300 and preventing transcriptional activation. ,PTM:In normoxia , is probably hydroxylated on Pro-405 and Pro-531 by EGLN1/PHD1 , EGLN2/PHD2 and/or EGLN3/PHD3. The hydroxylated prolines promote interaction with VHL , initiating rapid ubiquitination and subsequent proteasomal degradation. Under hypoxia , proline hydroxylation is impaired and ubiquitination is attenuated , resulting in stabilization. ,PTM:Phosphorylated on multiple sites in the CTAD. ,PTM:The iron and 2-oxoglutarate dependent 3-hydroxylation of asparagine is (S) stereospecific within HIF CTAD domains. ,similarity:Contains 1 basic helix-loop-helix (bHLH) domain. ,similarity:Contains 1 PAC (PAS-associated C-terminal) domain. ,similarity:Contains 2 PAS (PER-ARNT-SIM) domains. ,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Heterodimerizes with ARNT. Interacts with CREBBP. ,tissue specificity:Expressed in most tissues , with highest levels in placenta , lung and heart. Selectively expressed in endothelial cells. ,

| Validation Data

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: http://www.immunoway.com.cn
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
Acetyl-EPAS-1
(Lys385) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)