

Smad2/3 (Acetyl Lys19) Rabbit pAb

CatalogNo: YK0095

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- IHC, IF, WB

MW

- 60kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:50-200

WB 1:500-2000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human Smad2/3 (Acetyl-Lys19)

Specificity This antibody detects endogenous acetyl levels of Smad2/3 (Acetyl-Lys19) at Human:K19, Mouse:K19, Rat:K19. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):LGWkK

Target Information

Gene name Smad2/3 (Acetyl-Lys19)

Protein Name Smad2/3 (Acetyl-Lys19)

Organism	Gene ID	UniProt ID
Human	4088 ; 4087 ;	Q15796 ; P84022 ;

Cellular Localization


Cytoplasm . Nucleus . Cytoplasmic and nuclear in the absence of TGF-beta. On TGF-beta stimulation , migrates to the nucleus when complexed with SMAD4 (PubMed:9865696 , PubMed:21145499) . On dephosphorylation by phosphatase PPM1A , released from the SMAD2/SMAD4 complex , and exported out of the nucleus by interaction with RANBP1 (PubMed:16751101 , PubMed:19289081) . Localized mainly to the nucleus in the early stages of embryo development with expression becoming evident in the cytoplasm at the blastocyst and epiblast stages (By similarity) . .

Tissue specificity Expressed at high levels in skeletal muscle , endothelial cells , heart and placenta.

Function

Disease:Defects in SMAD2 are found in sporadic cases of colorectal carcinoma. ,Function:Transcriptional modulator activated by TGF-beta and activin type 1 receptor kinase. SMAD2 is a receptor-regulated SMAD (R-SMAD) . May act as a tumor suppressor in colorectal carcinoma. ,PTM:Acetylated on Lys-19 by coactivators in response to TGF-beta signaling , which increases transcriptional activity. Isoform short: Acetylation increases DNA binding activity in vitro and enhances its association with target promoters in vivo. ,PTM:In response to TGF-beta , ubiquitinated by NEDD4L; which promotes its degradation. ,PTM:Phosphorylated on one or several of Thr-220 , Ser-245 , Ser-250 , and Ser-255. In response to TGF-beta , phosphorylated on Ser-465/467 by TGF-beta and activin type 1 receptor kinases. Able to interact with SMURF2 when phosphorylated on Ser-465/467 , recruiting other proteins , such as SNON , for degradation. In response to decorin , the naturally occurring inhibitor of TGF-beta signaling , phosphorylated on Ser-240 by CaMK2. Phosphorylated by MAPK3 upon EGF stimulation; which increases transcriptional activity and stability , and is blocked by calmodulin. ,similarity:Belongs to the dwarfin/SMAD family. ,similarity:Contains 1 MH1 (MAD homology 1) domain. ,similarity:Contains 1 MH2 (MAD homology 2) domain. ,subcellular location:Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4. ,subunit:Found in a complex with SMAD3 and TRIM33 upon addition of TGF-beta. Interacts with SMAD3 and TRIM33. Interacts with SARA (SMAD anchor for receptor activation) ; may form trimers with the SMAD4 co-SMAD. Interacts with FOXH1 , homeobox protein TGIF , PEBP2-alpha subunit , CREB-binding protein (CBP) , EP300 and SKI. Interacts with SNON; when phosphorylated at Ser-465/467. Interacts (via PY-motif) with SMURF2. Interacts with AIP1 and HGS. Interacts with NEDD4L in response to TGF-beta (By similarity) . Interacts with LBXCOR1 and CORL2. ,tissue specificity:Expressed at high levels in skeletal muscle , heart and placenta. ,

Validation Data



Immunohistochemical analysis of paraffin-embedded human brain tumor. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min).

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Smad2/3 (Acetyl Lys19) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)