

JMJD2A Mouse mAb

CatalogNo: YM0389

| Key Features

Host Species

- Mouse

Reactivity

- Human

Applications

- WB,IHC,IF,ELISA

MW

- 121kD (Calculated)

| Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:200-1:1000

IF 1:200-1:1000

ELISA 1:10000

Not yet tested in other applications.

| Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

| Basic Information

Clonality Monoclonal

| Immunogen Information


Immunogen Purified recombinant fragment of human JMJD2A expressed in E. Coli.

Specificity JMJD2A Monoclonal Antibody detects endogenous levels of JMJD2A protein.


| Target Information

Gene name	KDM4A		
Protein Name	Lysine-specific demethylase 4A		
	Organism	Gene ID	UniProt ID
	Human	9682 ;	O75164 ;
	Mouse		Q8BW72 ;
Cellular Localization	Nucleus .		
Tissue specificity	Ubiquitous.		
Function	cofactor: Binds 1 Fe(2+) ion per subunit., Domain: The 2 Tudor domains recognize and bind methylated histone H3 'Lys-4' residue. Double Tudor domain has an interdigitated structure and the unusual fold is required for its ability to bind methylated histone tails. Trimethylated H3 'Lys-4' is bound in a cage of 3 aromatic residues, 2 of which are from the Tudor domain 2, while the binding specificity is determined by side-chain interactions involving residues from the Tudor domain 1. The Tudor domains are able to bind trimethylated histone H3 'Lys-4', trimethylated histone H3 'Lys-9', di- and trimethylated H4 'Lys-20'. , Function: Histone demethylase that specifically demethylates 'Lys-9' and 'Lys-36' residues of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-4', H3 'Lys-27' nor H4 'Lys-20'. Demethylates trimethylated H3 'Lys-9' and H3 'Lys-36' residue, while it has no activity on mono- and dimethylated residues. Demethylation of Lys residue generates formaldehyde and succinate. Participates in transcriptional repression of ASCL2 and E2F-responsive promoters via the recruitment of histone deacetylases and NCOR1, respectively., similarity: Belongs to the JHDM3 histone demethylase family., similarity: Contains 1 JmjC domain., similarity: Contains 1 JmjN domain., similarity: Contains 2 PHD-type zinc fingers., similarity: Contains 2 Tudor domains., subunit: Interacts with histone deacetylase proteins HDAC1, HDAC2 and HDAC3. Interacts with RB and NCOR1. Interacts with HTLV-1 Tax protein., tissue specificity: Ubiquitous.,		


| Validation Data



Western Blot analysis using JMJD2A Monoclonal Antibody against HEK293 (1) and JMJD2A-hlgGfc transfected HEK293 (2) cell lysate.



Immunohistochemistry analysis of paraffin-embedded colon cancer tissues (left) and human larynx cancer tissues (right) with DAB staining using JMJD2A Monoclonal Antibody.



Immunofluorescence analysis of NTERA-2 cells using JMJD2A Monoclonal Antibody (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
JMJD2A Mouse mAb

