

#### www.immunoway.com.cn

# NKX3.1 Mouse mAb

CatalogNo: YM0477

#### Key Features

Host Species

Mouse

Reactivity

Human

Applications • WB,IHC,IF,FC,ELISA

MW • 26kD (Calculated)

#### **Recommended Dilution Ratios**

WB 1:500-1:2000 IHC 1:200-1:1000 Flow Cyt 1:200-1:400 ELISA 1:10000 IF 1:50-200

#### **Storage**

Storage\*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### **Basic Information**

Clonality Monoclonal


### Immunogen Information

ImmunogenPurified recombinant fragment of human Nkx-3.1 expressed in E. Coli.SpecificityNkx-3.1 Monoclonal Antibody detects endogenous levels of Nkx-3.1 protein.


# Target Information

Gene name	NKX3-1 NKX3.1 NKX3A		
Protein Name	Homeobox protein Nkx-3.1 <b>Organism</b>	Gene ID	UniProt ID
	Human	<u>4824;</u>	<u>Q99801;</u>
	Mouse		<u>P97436;</u>
Cellular Localization	Nucleus .		
Tissue specificity	Highly expressed in the prostate and, at a lower level, in the testis.		
Function	Alternative products:Additional isoforms seem to exist,Disease:NKX3-1 has been thought to be one of the target gene of the 8p21 loss of heterozygosity, common in prostate cancer, but neither disruption of the coding region of the gene, nor mutations have been found in prostate cancer.,Function:Transcription factor, which binds preferentially the consensus sequence 5'-TAAGT[AG]-3' and can behave as a transcriptional repressor. Could play an important role in regulating proliferation of glandular epithelium and in the formation of ducts in prostate.,induction:By androgens and, in the LNCAP cell line, by estrogens. Androgenic control may be lost in prostate cancer cells during tumor progression from an androgen-dependent to an androgen-independent phase.,similarity:Belongs to the NK-3 homeobox family.,similarity:Contains 1 homeobox DNA-binding domain.,subunit:Interacts with serum response factor (SRF) (By similarity). Interacts with SPDEF. Interacts with WDR77.,tissue specificity:Highly expressed in the prostate and, at a lower level, in the testis.,		


# Validation Data



Western Blot analysis using Nkx-3.1 Monoclonal Antibody against LNCaP (1) cell lysate.



Immunohistochemistry analysis of paraffin-embedded human prostata tissues (A, B) with DAB staining using Nkx-3.1 Monoclonal Antibody.



Flow cytometric analysis of PC-3 cells using Nkx-3.1 Monoclonal Antibody (green) and negative control (purple).

# **Contact information**

Orders:	order.cn@immunoway.com
Support:	support.cn@immunoway.com
Telephone:	400-8787-807(China)
Website:	http://www.immunoway.com.cn
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **NKX3.1 Mouse mAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents