

### TrxR2 Mouse mAb

CatalogNo: YM1332

# **Key Features**

**Host Species** 

Reactivity
• Human

**Applications** 

WB

Mouse

MW56kD (Observed)

### **Recommended Dilution Ratios**

WB 1:1000

## Storage

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

# **Basic Information**

**Clonality** Monoclonal

### Immunogen Information

**Immunogen** Purified recombinant human TXNRD2 protein fragments expressed in E.coli.

**Specificity** This antibody detects endogenous levels of TXNRD2 and does not cross-react with

related proteins.

# | Target Information

Gene name TXNRD2

#### **Protein Name**

Thioredoxin reductase 2 mitochondrial

Organism	Gene ID	UniProt ID
Human	<u>10587;</u>	Q9NNW7;

### Cellular Localization


Mitochondrion.

**Tissue specificity** Highly expressed in the prostate, ovary, liver, testis, uterus, colon and small intestine. Intermediate levels in brain, skeletal muscle, heart and spleen. Low levels in placenta, pancreas, thymus and peripheral blood leukocytes. According to PubMed:10608886, high levels in kidney, whereas according to PubMed:9923614, levels are low. High expression is observed in the adrenal cortex (PubMed:24601690).

#### **Function**

Catalytic activity: Thioredoxin + NADP(+) = thioredoxin disulfide +NADPH., cofactor: FAD., Function: Maintains thioredoxin in a reduced state. Implicated in the defenses against oxidative stress. May play a role in redox-regulated cell signaling,,miscellaneous:The active site is a redox-active disulfide bond. The selenocysteine residue is essential for enzymatic activity., sequence Caution: Translated as Sec., similarity: Belongs to the class-I pyridine nucleotide-disulfide oxidoreductase family., subunit: Homodimer., tissue specificity: Highly expressed in the prostate, ovary, liver, testis, uterus, colon and small intestine. Intermediate levels in brain, skeletal muscle, heart and spleen. Low levels in placenta, pancreas, thymus and peripheral blood leukocytes. According to PubMed:10608886, high levels in kidney, whereas according to PubMed:9923614 levels are low.,

### Validation Data



Western blot detection of TXNRD2 in A431,U251,U2OS,Lncap and Hela cell lysates and using TXNRD2 mouse mAb (1:1000 diluted). Predicted band size: 56KDa.Observed band size: 56KDa.

# I Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

400-8787-807(China) Telephone:

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

TrxR2 Mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents