

www.immunoway.com.cn

DDX3 Mouse mAb

CatalogNo: YM1415 Comparable Abs

Key Features

Host SpeciesReactivity• Mouse• Human,Rat,Mouse,Monkey

ApplicationsWB,IP,ICC

MW • 75kD (Observed)

Recommended Dilution Ratios

WB 1:1000 ICC 1:200

Storage

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Monoclonal

Immunogen Information

proteins.

ddx3x


ImmunogenPurified recombinant human DDX3 protein fragments expressed in E.coli.SpecificityThis antibody detects endogenous levels of DDX3 and does not cross-react with related

Target Information


Gene name

Protein Name	Organism	Gene ID	UniProt ID
	Human	<u>1654;</u>	<u>000571;</u>
	Mouse		<u>Q62167;</u>
Cellular Localization	Cell membrane . Nucleus . Cytoplasm . Cytoplasm, Stress granule . Inflammasome . Cell projection, lamellipodium . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Shuttles between the nucleus and the cytosol (PubMed:15507209, PubMed:18636090, PubMed:29899501, PubMed:31575075, PubMed:30131165). Exported from the nucleus partly through the XPO1/CRM1 system and partly through NXF1/TAP (PubMed:15507209, PubMed:18636090, PubMed:18636090, PubMed:38596238, PubMed:31575075, PubMed:30131165). Localizes to nuclear pores on the outer side of the nuclear membrane (PubMed:15507209). In the cytosol, partly colocalizes with mitochondria (PubMed:20127681). At G0, predominantly located in nucleus. In G1/S phase, predominantly cytoplasmic (PubMed:22034099). During prophase/prometaphase, localizes in close proximity to the condensing chromosomes (PubMed:30131165, PubMed:21730191). During telophase, localizes around the newly synthesized nuclear membrane and in the cytoplasm (PubMed:22034099). Colocalizes with TRPV4 at the plasma membrane. When TRPV4 channel is activated, intracellular Ca(2+) levels increase and the calmodulin/CAMKII pathway is activated, relocalizes to the nucleus (PubMed:23413191). At the leading edge of migrating fibroblasts, colocalizes with CAPRIN1 and PABPC1 (PubMed:28733330). Localizes to centrosome throughout the cell cycle and associates with TP53 at centrosome during mitosis (PubMed:28842590). Translocates to the nucleus in response to HPIV-3 virus-mediated infection (PubMed:31575075)		
Tissue specificity	Widely expressed (PubMed:15294 (PubMed:15294876). Expressed in PubMed:16301996).		sed in spermatids at protein level) (PubMed:16818630,
Function	of incompletely spliced HIV-1 Rev specifically with hepatitis C virus of location.,similarity:Belongs to the box helicase family. DDX3/DED1 s domain.,similarity:Contains 1 helic	RNAs. Also involved in core protein resulting i DEAD box helicase far ubfamily.,similarity:Co case C-terminal domai and, at low levels, the plexes (NPC). Shuttles nanner.,subunit:Found	in a change in intracellular mily.,similarity:Belongs to the DEAD ontains 1 helicase ATP-binding n.,subcellular location:Located roughout the cytoplasm. Located to s between the nucleus and the l in a complex with Rev and XPO1.


Validation Data



Western blot detection of DDX3 in Hela,3T3,C6,COS7,K562 and Jurkat cell lysate using DDX3 mouse mAb (1:1000 diluted).Predicted band size: 75KDa.Observed band size: 75KDa.



Immunocytochemistry staining of HeLa cells fixed with 4% Paraformaldehyde and using DDX3 mouse mAb (dilution 1:200).



Immunoprecipitation analysis of Hela cell lysates using DDX3 mouse mAb.

Contact information

Orders:	order.cn@immunoway.com
Support:	support.cn@immunoway.com
Telephone:	400-8787-807(China)
Website:	http://www.immunoway.com.cn
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **DDX3 Mouse mAb**