

## Caspase-1 (PTR1290) Mouse mAb

CatalogNo: YM4685

### | Key Features

#### Host Species

- Mouse

#### Reactivity

- Human, Mouse, Rat, Horse, Pig,

#### Applications

- WB, IF, ELISA

#### MW

- 45kD (Calculated)  
45kD (Observed)

#### Isotype

- IgG2a, Kappa

### | Recommended Dilution Ratios

**WB 1:500-2000**

**IF 1:100-500**

**ELISA 1:1000-5000**

### | Storage

**Storage\*** -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

### | Basic Information

**Clonality** Monoclonal

**Clone Number** PTR1290

### | Immunogen Information

**Immunogen** Synthesized peptide derived from human Caspase-1 AA range: 350-404

**Specificity** This antibody detects endogenous levels of Caspase-1 protein.

## Target Information

**Gene name** CASP1 IL1BC IL1BCE

**Protein Name** Caspase-1


Organism	Gene ID	UniProt ID
Human	<a href="#">834</a> ;	<a href="#">P29466</a> ;
Mouse	<a href="#">12362</a> ;	<a href="#">P29452</a> ;
Rat		<a href="#">P43527</a> ;

**Cellular Localization** Cytoplasm

**Tissue specificity** Expressed in larger amounts in spleen and lung. Detected in liver, heart, small intestine, colon, thymus, prostate, skeletal muscle, peripheral blood leukocytes, kidney and testis. No expression in the brain.

**Function** Thiol protease involved in a variety of inflammatory processes by proteolytically cleaving other proteins, such as the precursors of the inflammatory cytokines interleukin-1 beta (IL1B) and interleukin 18 (IL18) as well as the pyroptosis inducer Gasdermin-D (GSDMD), into active mature peptides . Plays a key role in cell immunity as an inflammatory response initiator: once activated through formation of an inflammasome complex, it initiates a pro-inflammatory response through the cleavage of the two inflammatory cytokines IL1B and IL18, releasing the mature cytokines which are involved in a variety of inflammatory processes . Cleaves a tetrapeptide after an Asp residue at position P1 . Also initiates pyroptosis, a programmed lytic cell death pathway, through cleavage of GSDMD . In contrast to cleavage of interleukins IL1B and IL1B, recognition and cleavage of GSDMD is not strictly dependent on the consensus cleavage site but depends on an exosite interface on CASP1 that recognizes and binds the Gasdermin-D, C-terminal (GSDMD-CT) part . Upon inflammasome activation, during DNA virus infection but not RNA virus challenge, controls antiviral immunity through the cleavage of CGAS, rendering it inactive . In apoptotic cells, cleaves SPHK2 which is released from cells and remains enzymatically active extracellularly . ; [Isoform Delta]: Apoptosis inactive. ; [Isoform Epsilon]: Apoptosis inactive.

## Validation Data



Whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with anti-Caspase-1 (PTR1290) antibody. The HRP-conjugated Goat anti-Mouse IgG(H + L) antibody was used to detect the antibody. Lane 1: THP-1

## | Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:

**Caspase-1  
(PTR1290) Mouse  
mAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)