

COX-2 (ABT-COX2) Mouse mAb

CatalogNo: YM4881

Key Features

Host Species

- Mouse

Reactivity

- Human

Applications

- IHC, WB, IF, ELISA

MW

- 69kD (Calculated)
- 70kD (Observed)

Isotype

- IgG2b, Kappa

Recommended Dilution Ratios

IHC 1:200-1000**WB 1:500-2000****IF 1:100-500****ELISA 1:1000-5000**

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Basic Information

Clonality Monoclonal**Clone Number** ABT-COX2


Immunogen Information

Immunogen Synthesized peptide derived from human COX-2 (cyclooxygenase-2) AA range: 500-604**Specificity** The antibody can specifically recognize human COX-2 protein. In western blotting of Hela and A549 cell lysates, the antibody can label a 70 kDa band corresponding to COX-2.


| Target Information

Gene name	PTGS2 COX2		
Protein Name	Prostaglandin G/H synthase 2 (Cyclooxygenase-2) (COX-2) (PHS II) (Prostaglandin H2 synthase 2) (PGH synthase 2) (PGHS-2) (Prostaglandin-endoperoxide synthase 2)		
	Organism	Gene ID	UniProt ID
	Human	5743 ;	P35354 ;
Cellular Localization	Cytoplasmic		
Tissue specificity	Endothelial cell,Epidermal keratinocytes in primary culture,Lung,Pe		
Function	Catalytic activity:Arachidonate + AH(2) + 2 O(2) = prostaglandin H(2) + A + H(2)O.,cofactor:Binds 1 heme B (iron-protoporphyrin IX) group per subunit.,Disease:Likely to play a role in inflammatory diseases such as rheumatoid arthritis.,Function:May have a role as a major mediator of inflammation and/or a role for prostanoid signaling in activity-dependent plasticity.,induction:By cytokines and mitogens.,miscellaneous:This enzyme acts both as a dioxygenase and as a peroxidase.,miscellaneous:This enzyme is the target of nonsteroidal anti-inflammatory drugs such as aspirin.,pathway:Lipid metabolism; prostaglandin biosynthesis.,similarity:Belongs to the prostaglandin G/H synthase family.,similarity:Contains 1 EGF-like domain.,subunit:Homodimer.,		


| Validation Data



Human colon carcinoma tissue was stained with Anti-COX-2 (ABT-COX2) Antibody



Various whole cell lysates were separated by 8% SDS-PAGE, and the membrane was blotted with anti-COX-2. The HRP-conjugated anti-Mouse IgG antibody was used to detect the antibody.



Whole cell lysates of HeLa were separated by 10% SDS-PAGE, and the membrane was blotted with anti-COX-2 antibody. The HRP-conjugated anti-Mouse IgG antibody was used to detect the antibody.

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
COX-2 (ABT-COX2)
Mouse mAb