

PIK3CA H1047R (PT0281R) Rabbit mAb (Ready to Use)

CatalogNo: YM7195R **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human,

Applications

- IHC

Isotype

- IgG1, Kappa

Recommended Dilution Ratios

Ready to use for IHC

Storage

Storage* 2°C to 8°C/1 year

Formulation The prediluted ready-to-use antibody is diluted in phosphate buffer saline containing stabilizing protein and 0.05% Proclin 300

Basic Information

Clonality Monoclonal

Clone Number PT0281R

Immunogen Information

Immunogen Synthesized peptide derived from human PIK3CA H1047R AA range:1000-1068

Specificity This antibody detects endogenous levels of PIK3CA H1047R

Target Information

Gene name	PIK3CA		
Protein Name	5-bisphosphate 3-kinase 110 kDa catalytic subunit alpha;5-bisphosphate 3-kinase catalytic subunit alpha isoform;caPI3K;CLOVE;CWS5;MCAP;MCM;MCMTc;MGC142161;MGC142163;p110 alpha;p110alpha;Phosphatidylinositol 3 kinase catalytic alpha polypeptide;Phosphatidylinositol 3 kinase catalytic 110 KD alpha;Phosphatidylinositol 4 5 bisphosphate 3 kinase catalytic subunit alpha;Phosphatidylinositol 4 5 bisphosphate 3 kinase catalytic subunit alpha isoform;Phosphatidylinositol 4,5 bisphosphate 3 kinase 110 kDa catalytic subunit alpha;Phosphatidylinositol-4;Phosphoinositide 3 kinase catalytic alpha polypeptide;PI 3 Kinase catalytic subunit alpha;PI3 kinase p110 subunit alpha;PI3-kinase subunit alpha;PI3K;PI3K-alpha;PI3KC A;PIK3C A;Pik3ca;PK3CA;PK3CA_HUMAN;PtdIns 3 kinase p110;PtdIns-3-kinase subunit alpha;PtdIns-3-kinase subunit p110-alpha;Serine/threonine protein kinase PIK3CA		
	Organism	Gene ID	UniProt ID
	Human	5290 ;	P42336 ;
	Mouse		P42337 ;
Cellular Localization	Cytoplasmic		
Tissue specificity	Gastric adenocarcinoma with PIK3CA H1047R protein expression		
Function	Catalytic activity:ATP + 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate = ADP + 1-phosphatidyl-1D-myo-inositol 3,4,5-trisphosphate.,Disease:Defects in PIK3CA are associated with breast cancer [MIM:114480].,Disease:Defects in PIK3CA are associated with colorectal cancer (CRC) [MIM:114500].,Disease:Defects in PIK3CA are associated with ovarian cancer [MIM:167000]. Ovarian cancer is the leading cause of death from gynecologic malignancy. It is characterized by advanced presentation with loco-regional dissemination in the peritoneal cavity and the rare incidence of visceral metastases. These typical features relate to the biology of the disease, which is a principal determinant of outcome.,Disease:Defects in PIK3CA may underlie hepatocellular carcinoma (HCC) [MIM:114550].,Disease:PI3KCA mutations affecting exons 9 and 20 display gender-and tissue-specific patterns, thus suggesting that the different amino acid changes could exert distinct functional effects on the oncogenic properties of this enzyme. Furthermore, sexual dimorphisms and tissue specific factors might directly or indirectly influence the occurrence of PI3KCA cancer alleles.,Function:Phosphorylates PtdIns, PtdIns4P and PtdIns(4,5)P2 with a preference for PtdIns(4,5)P2.,similarity:Belongs to the PI3/PI4-kinase family.,similarity:Contains 1 C2 domain.,similarity:Contains 1 PI3K/PI4K domain.,subunit:Heterodimer of a p110 (catalytic) and a p85 (regulatory) subunit. Binds to IRS1 in nuclear extracts. Interacts with RUFY3.,		

| Validation Data

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**PIK3CA H1047R
(PT0281R) Rabbit
mAb (Ready to Use)**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)