

NOX2 (PT0125R) PT™ Rabbit mAb

CatalogNo: YM8068 Recombinant R

Key Features

Host Species

Rabbit

MW
• 65kD (Calculated)

65kD (Observed)

Reactivity

Human, Mouse, Rat

Isotype

IgG,Kappa

ApplicationsWB,ELISA

Recommended Dilution Ratios

WB 1:1000-5000 ELISA 1:5000-20000

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Basic Information

Clonality Monoclonal

Clone Number PT0125R

Immunogen Information

Specificity Endogenous

| Target Information

Gene name

CYBB

Protein Name

Cytochrome b-245 heavy chain

Organism	Gene ID	UniProt ID	
Human	<u>1536</u> ;	<u>P04839</u> ;	
Mouse		<u>Q61093;</u>	

Cellular Localization


Membranous

Tissue specificity Detected in neutrophils (at protein level).


Function

cofactor:FAD.,Disease:Defects in CYBB are a cause of chronic granulomatous disease X-linked (XCGD) [MIM:306400]. Chronic granulomatous disease is a genetically heterogeneous disorder characterized by the inability of neutrophils and phagocytes to kill microbes that they have ingested. Patients suffer from life-threatening bacterial/fungal infections.,Function:Critical component of the membrane-bound oxidase of phagocytes that generates superoxide. It is the terminal component of a respiratory chain that transfers single electrons from cytoplasmic NADPH across the plasma membrane to molecular oxygen on the exterior. Also functions as a voltage-gated proton channel that mediates the H(+) currents of resting phagocytes. It participates in the regulation of cellular pH and is blocked by zinc.,online information:CYBB deficiency database,PTM:Glycosylated.,similarity:Contains 1 FAD-binding FR-type domain.,similarity:Contains 1 ferric oxidoreductase domain.,subunit:Composed of a heavy chain (beta) and a light chain (alpha).,


I Validation Data




Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-NOX2 antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: HepG2 Lane 2: Jurkat Lane 3: Mouse Brain Predicted band size: 65kDa Observed band size: 65kDa



Western blot analysis of lysate from K562 cells, using CYBB Antibody.



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the primary antibody was used at 4°C, over night with a 1:2500 dilution . The Dylight 800-conjugated Goat anti-Rabbit antibody(Cat:RS23920) was used to detect the antibody. Lane1: HL60 - Human promyelocytic leukemia cell Lane2: HUVEC - Human umbilical vein endothelial cell Lane3: Human CAF - Human cancer-associated fibroblast Lane4: HCT116 - Human colorectal carcinoma Lane5: A549 - Human lung carcinoma Lane6: Jurkat - Human T lymphocyte leukemia Lane7: HT22 - Mouse hippocampal neuronal Lane8: Thp-1 - Human monocytic leukemia Lane9: HK-2 - Human proximal tubular epithelial Lane10: PC-12 - Rat adrenal pheochromocytoma Lane11: 3T3-L1 - Mouse embryonic fibroblast Lane12: PC-3 - Human prostate adenocarcinoma Lane13: U-14 - Mouse cervical carcinoma Lane14: 293F - HEK293 derivative, adapted for suspension culture Predicted band size: 65kDa Observed band size: 65kDa



Immunofluorescence analysis of MCF7 cell. 1,primary Antibody was diluted at $1:100(4 \sim C \text{ overnight})$. 2, Goat Anti Rabbit IgG (H&L) - AFluor 594 Secondary antibody(catalog No: RS3611) was diluted at 1:500(room temperature, 50min).

I Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
NOX2 (PT0125R)
PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents