

## Keap1 (PT0444R) PT™ Rabbit mAb

CatalogNo: YM8283 **Recombinant** 

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, IHC, IF, ELISA

#### MW

- 70kD (Calculated)  
60-70kD (Observed)

#### Isotype

- IgG, Kappa

### Recommended Dilution Ratios

IHC 1:200-1:1000

WB 1:2000-1:10000

IF 1:200-1:1000

ELISA 1:5000-1:20000

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

### Basic Information

**Clonality** Monoclonal

**Clone Number** PT0444R


### Immunogen Information

**Specificity** Endogenous


## | Target Information

Gene name	KEAP1 INRF2 KIAA0132 KLHL19		
Protein Name	Kelch-like ECH-associated protein 1 (Cytosolic inhibitor of Nrf2) (INrf2) (Kelch-like protein 19)		
	Organism	Gene ID	UniProt ID
	Human	<a href="#">9817</a> ;	<a href="#">Q14145</a> ;
	Mouse	<a href="#">50868</a> ;	<a href="#">Q9Z2X8</a> ;
	Rat	<a href="#">117519</a> ;	<a href="#">P57790</a> ;
Cellular Localization	Cytoplasm, Nucleus		
Tissue specificity	Broadly expressed, with highest levels in skeletal muscle.		
Function	Disease:Defects in KEAP1 may be a cause of breast cancer.,Disease:Defects in KEAP1 may be involved in non small cell lung carcinomas (NSCLC) and lung adenocarcinoma.,Domain:The Kelch repeats mediate interaction with NF2L2/NRF2, BPTF and PGAM5.,enzyme regulation:Ubiquitination and subsequent degradation of PGAM5 is inhibited by oxidative stress and sulforaphane.,Function:Retains NFE2L2/NRF2 in the cytosol. Functions as substrate adapter protein for the E3 ubiquitin ligase complex formed by CUL3 and RBX1. Targets NFE2L2/NRF2 for ubiquitination and degradation by the proteasome, thus resulting in the suppression of its transcriptional activity and the repression of antioxidant response element-mediated detoxifying enzyme gene expression. May also retain BPTF in the cytosol. Targets PGAM5 for ubiquitination and degradation by the proteasome.,PTM:Ubiquitinated and subject to proteasomal degradation.,similarity:Contains 1 BACK (BTB/Kelch associated) domain.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 6 Kelch repeats.,subcellular location:Shuttles between cytoplasm and nucleus.,subunit:Homodimer. Interacts with the N-terminal regulatory domain of NF2L2/NRF2. Interacts with BPTF and PTMA. Interacts with CUL3. Part of a complex that contains KEAP1, CUL3 and RBX1. Interacts with PGAM5.,tissue specificity:Broadly expressed, with highest levels in skeletal muscle.,		


## | Validation Data




Mouse brain was stained with anti-Keap1 rabbit antibody




Rat brain was stained with anti-Keap1 rabbit antibody




Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-Keap1 antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: HEK293 Lane 2: RAW264.7 Lane 3: Rat spleen Predicted band size: 70kDa Observed band size: 60-70kDa




Human brain was stained with anti-Keap1 rabbit antibody



Human prostate was stained with anti-Keap1 rabbit antibody



Mouse kidney was stained with anti-Keap1 Rabbit antibody



Mouse kidney was stained with anti-Keap1 Rabbit antibody

## | Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**Keap1 (PT0444R)**  
**PT™ Rabbit mAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)