

ATM (Phospho Ser1981) (PT0902R) PT™ Rabbit mAb

CatalogNo: YM8671 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, IP, ELISA, CHIP, Cut&Tag

MW

- 351kD
(Calculated)
351kD
(Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Recommended Dilution Ratios

IHC 1:200-1:1000

WB 1:10000-1:50000

IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

CHIP 1:50-1:100

Cut&Tag 1:50-1:100

Basic Information

Clonality Monoclonal

Clone Number PT0902R

Immunogen Information

Immunogen The specific immunogen used to produce this antibody is proprietary information.

Specificity ATM (Phospho Ser1981) Monoclonal Antibody detects endogenous levels of Phospho Atm around the phosphorylation site of S1981(human), S1896(mouse), S1927(rat) protein. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):EGsQS


| **Target Information**

Gene name	ATM		
Protein Name	Serine-protein kinase ATM		
	Organism	Gene ID	UniProt ID
	Human	472;	Q13315;
	Mouse	11920;	Q62388;
Cellular Localization	Nucleus . Cytoplasmic vesicle . Cytoplasm , cytoskeleton , microtubule organizing center , centrosome . Primarily nuclear. Found also in endocytic vesicles in association with beta-adaptin. .		
Tissue specificity	Found in pancreas , kidney , skeletal muscle , liver , lung , placenta , brain , heart , spleen , thymus , testis , ovary , small intestine , colon and leukocytes.		


Function

Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,Disease:Defects in ATM are the cause of ataxia telangiectasia (AT) [MIM:208900]; also known as Louis-Bar syndrome , which includes four complementation groups: A , C , D and E. This rare recessive disorder is characterized by progressive cerebellar ataxia , dilation of the blood vessels in the conjunctiva and eyeballs , immunodeficiency , growth retardation and sexual immaturity. AT patients have a strong predisposition to cancer; about 30% of patients develop tumors , particularly lymphomas and leukemias. Cells from affected individuals are highly sensitive to damage by ionizing radiation and resistant to inhibition of DNA synthesis following irradiation. ,Disease:Defects in ATM contribute to B-cell chronic lymphocytic leukemia (BCLL) . BCLL is the commonest form of leukemia in the elderly. It is characterized by the accumulation of mature CD5+ B lymphocytes , lymphadenopathy , immunodeficiency and bone marrow failure. ,Disease:Defects in ATM contribute to B-cell non-Hodgkin lymphomas (BNHL) , including mantle cell lymphoma (MCL) . ,Disease:Defects in ATM contribute to T-cell acute lymphoblastic leukemia (TALL) and T-prolymphocytic leukemia (TPLL) . TPLL is characterized by a high white blood cell count , with a predominance of prolymphocytes , marked splenomegaly , lymphadenopathy , skin lesions and serous effusion. The clinical course is highly aggressive , with poor response to chemotherapy and short survival time. TPLL occurs both in adults as a sporadic disease and in younger AT patients. ,Domain:The FATC domain is required for interaction with HTATIP. ,enzyme regulation:Inhibited by wortmannin. ,Function:Serine/threonine protein kinase which activates checkpoint signaling upon double strand breaks (DSBs) , apoptosis and genotoxic stresses such as ionizing ultraviolet A light (UVA) , thereby acting as a DNA damage sensor. Recognizes the substrate consensus sequence [ST]-Q. Phosphorylates 'Ser-139' of histone variant H2AX/H2AFX at double strand breaks (DSBs) , thereby regulating DNA damage response mechanism. Also involved in signal transduction and cell cycle control. May function as a tumor suppressor. Necessary for activation of ABL1 and SAPK. Phosphorylates p53/TP53 , FANCD2 , NFKBIA , BRCA1 , CTIP , nibrin (NBN) , TERF1 , RAD9 and DCLRE1C. May play a role in vesicle and/or protein transport. Could play a role in T-cell development , gonad and neurological function. ,induction:By ionizing radiation. ,online information:Ataxia telangiectasia mutated entry ,PTM:Acetylation , on DNA damage , is required for activation of the kinase activity , dimer-monomer transition , and subsequent autophosphorylation on Ser-1981. Acetylated in vitro by HTATIP/TIP60. ,PTM:Phosphorylated by NUA1/ARK5. Autophosphorylation on Ser-367 , Ser-1983 , Ser-1981 correlates with DNA damage-mediated activation of the kinase. ,similarity:Belongs to the PI3/PI4-kinase family. ATM subfamily. ,similarity:Contains 1 FAT domain. ,similarity:Contains 1 FATC domain. ,similarity:Contains 1 PI3K/PI4K domain. ,subcellular location:Primarily nuclear. Found also in endocytic vesicles in association with beta-adaptin. ,subunit:Dimers or tetramers in inactive state. On DNA damage , autophosphorylation dissociates ATM into monomers rendering them catalytically active. Binds DNA ends , p53/TP53 , ABL1 , BRCA1 , NBN/nibrin and TERF1. Part of the BRCA1-associated genome surveillance complex (BASC) , which contains BRCA1 , MSH2 , MSH6 , MLH1 , ATM , BLM , PMS2 and the RAD50-MRE11-NBN protein complex. This association could be a dynamic process changing throughout the cell cycle and within subnuclear domains. DNA damage promotes association with RAD17. Interacts with EEF1E1; the interaction , induced on DNA damage , upregulates TP53. Interacts with DCLRE1C , MYST1 , HTATIP , OBFC2B , ATMIN and CEP164. Interacts with the beta-adaptin complex subunits , AP2B1 AND AP3B2; the interaction occurs in cytoplasmic vesicles. ,tissue specificity:Found in pancreas , kidney , skeletal muscle , liver , lung , placenta , brain , heart , spleen , thymus , testis , ovary , small intestine , colon and leukocytes. ,

| Validation Data



Various whole cell lysates were separated by 4-8% SDS-PAGE, and the membrane was blotted with anti-ATM (Phospho Ser1981) (PT0902R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: HeLa Lane 2: HeLa treated with UV for 1 hour, then recovered for 1 hour Predicted band size: 351kDa Observed band size: 351kDa



Human breast carcinoma was stained with anti-ATM (Phospho Ser1981) (PT0902R) Rabbit antibody

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
ATM (Phospho Ser1981) (PT0902R) PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)