

Rpb1 (Phospho Ser5) (PT0923R) PT™ Rabbit mAb

CatalogNo: YM8717 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, IP, ELISA

MW

- 217kD (Calculated)
217kD (Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

IHC 1:200-1:1000

WB 1:500-1:2000

IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

Basic Information

Clonality Monoclonal

Clone Number PT0923R

Immunogen Information

Immunogen The specific immunogen used to produce this antibody is proprietary information.

Specificity Rpb1 (Phospho Ser5) Monoclonal Antibody detects endogenous levels of Rpb1 protein only when phosphorylated at Ser5. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):GGgPP


| Target Information

Gene name	POLR2A POLR2		
Protein Name	Rpb1 CTD (Ser5)		
	Organism	Gene ID	UniProt ID
	Human	5430 ;	P24928 ;
	Mouse	20020 ;	P08775 ;
Cellular Localization	Nucleus . Cytoplasm . Chromosome . Hypophosphorylated form is mainly found in the cytoplasm , while the hyperphosphorylated and active form is nuclear (PubMed:26566685) . Co-localizes with kinase SRPK2 and helicase DDX23 at chromatin loci where unscheduled R-loops form (PubMed:28076779) . .		
Tissue specificity	Fetal pancreas ,Testis ,		


Function

Catalytic activity: Nucleoside triphosphate + RNA (n) = diphosphate + RNA (n+1) .
Function: DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Largest and catalytic component of RNA polymerase II which synthesizes mRNA precursors and many functional non-coding RNAs. Forms the polymerase active center together with the second largest subunit. Pol II is the central component of the basal RNA polymerase II transcription machinery. It is composed of mobile elements that move relative to each other. RPB1 is part of the core element with the central large cleft , the clamp element that moves to open and close the cleft and the jaws that are thought to grab the incoming DNA template. At the start of transcription , a single stranded DNA template strand of the promoter is positioned within the central active site cleft of Pol II. A bridging helix emanates from RPB1 and crosses the cleft near the catalytic site and is thought to promote translocation of Pol II by acting as a ratchet that moves the RNA-DNA hybrid through the active site by switching from straight to bent conformations at each step of nucleotide addition. During transcription elongation , Pol II moves on the template as the transcript elongates. Elongation is influenced by the phosphorylation status of the C-terminal domain (CTD) of Pol II largest subunit (RPB1) , which serves as a platform for assembly of factors that regulate transcription initiation , elongation , termination and mRNA processing. Acts as a RNA-dependent RNA polymerase when associated with small delta antigen of Hepatitis delta virus , acting both as a replicate and transcriptase for the viral RNA circular genome. ,**miscellaneous:** The binding of ribonucleoside triphosphate to the RNA polymerase II transcribing complex probably involves a two-step mechanism. The initial binding seems to occur at the entry (E) site and involves a magnesium ion temporarily coordinated by three conserved aspartate residues of the two largest RNA Pol II subunits. The ribonucleoside triphosphate is transferred by a rotation to the nucleotide addition (A) site for pairing with the template DNA. The catalytic A site involves three conserved aspartate residues of the RNA Pol II largest subunit which permanently coordinate a second magnesium ion. ,**PTM:** The tandem 7 residues repeats in the C-terminal domain (CTD) can be highly phosphorylated. The phosphorylation activates Pol II. Phosphorylation occurs mainly at residues 'Ser-2' and 'Ser-5' of the heptapeptide repeat. The phosphorylation state is believed to result from the balanced action of site-specific CTD kinases and phosphatases , and a "CTD code" that specifies the position of Pol II within the transcription cycle has been proposed. ,**similarity:** Belongs to the RNA polymerase beta' chain family. ,**similarity:** Contains 1 C2H2-type zinc finger. ,**subunit:** Component of the RNA polymerase II (Pol II) complex consisting of 12 subunits (By similarity) . The phosphorylated C-terminal domain interacts with FNPB3 and SYNERGIP. Interacts with SAFB/SAFB1. Interacts with CCNL1 and MYO1C (By similarity) . Interacts with CCNL2 and SFRS19. Component of a complex which is at least composed of HTATSF1/Tat-SF1 , the P-TEFb complex components CDK9 and CCNT1 , RNA polymerase II , SUPT5H , and NCL/nucleolin. Interacts with PAF1. ,

Validation Data



Various whole cell lysates were separated by 4-8% SDS-PAGE, and the membrane was blotted with anti-Rpb1 (Phospho Ser5) (PT0923R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: NIH-3T3 Lane 2: NIH-3T3 was treated with TSA (400nM) for 18 hours Lane 3: C6 Lane 4: C6 was treated with TSA (500ng/ml) for 4 hours Predicted band size: 217kDa Observed band size: 217kDa



Human ovarian carcinoma was stained with anti-Rpb1 (Phospho Ser5) (PT0923R) Rabbit antibody

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Rpb1 (Phospho Ser5) (PT0923R) PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)