

PLC γ 1 (Phospho Tyr783) (PT0758R) PT™ Rabbit mAb

CatalogNo: YM8720 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, IP, ELISA

MW

- 149kD (Calculated)
149kD (Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

IHC 1:200-1:1000

WB 1:2000-1:10000

IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

Basic Information

Clonality Monoclonal

Clone Number PT0758R

Immunogen Information


Immunogen The specific immunogen used to produce this antibody is proprietary information.

Specificity PLCγ1 (Phospho Tyr783) Monoclonal Antibody detects endogenous levels of PLC γ1 protein only when phosphorylated at Y783. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):GFyVE


| Target Information

Gene name	PLCG1		
Protein Name	1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1		
	Organism	Gene ID	UniProt ID
	Human	5335 ;	P19174 ;
	Mouse	18803 ;	Q62077 ;
	Rat	25738 ;	P10686 ;
Cellular Localization	Cell projection , lamellipodium . Cell projection , ruffle . Rapidly redistributed to ruffles and lamellipodia structures in response to epidermal growth factor (EGF) treatment. .		
Tissue specificity	Brain ,Epithelium ,Testis ,Vein ,		
Function	Catalytic activity:1-phosphatidyl-1D-myo-inositol 4 ,5-bisphosphate + H (2) O = 1D-myo-inositol 1 ,4 ,5-trisphosphate + diacylglycerol. ,cofactor:Calcium. ,Domain:The SH3 domain mediates interaction with CLNK (By similarity) . The SH3 domain also mediates interaction with RALGPS1. ,Function:PLC-gamma is a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor) -activated tyrosine kinase. ,PTM:The receptor-mediated activation of PLC-gamma-1 and PLC-gamma-2 involves their phosphorylation by tyrosine kinases in response to ligation of a variety of growth factor receptors and immune system receptors. ,PTM:Ubiquitinated by CBLB in activated T-cells. ,similarity:Contains 1 C2 domain. ,similarity:Contains 1 EF-hand domain. ,similarity:Contains 1 PH domain. ,similarity:Contains 1 PI-PLC X-box domain. ,similarity:Contains 1 PI-PLC Y-box domain. ,similarity:Contains 1 SH3 domain. ,similarity:Contains 2 PH domains. ,similarity:Contains 2 SH2 domains. ,similarity:Contains 3 PH domains. ,subunit:Interacts with AGAP2 via its SH3 domain (By similarity) . Interacts with phosphorylated LAT upon TCR activation. Interacts with the Pro-rich domain of TNK1 via its SH3 domain. Associates with BLNK , VAV1 , GRB2 and NCK1 in a B-cell antigen receptor-dependent fashion. Interacts with CBLB in activated T-cells; which inhibits phosphorylation. Interacts with SHB. Interacts via its SH3 domain with the Arg/Gly-rich-flanked Pro-rich domains of KHDRBS1/SAM68. This interaction is selectively regulated by arginine methylation of KHDRBS1/SAM68. Interacts with INPP5D/SHIP1 and CLNK (By similarity) . Interacts with RALGPS1. Interacts (via SH3 domain) with HEV ORF3 protein. ,		

| Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-PLCγ1 (Phospho Tyr783) (PT0758R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: HeLa Lane 2: HeLa was treated with Pervanadate (10mM) for 1 hour Lane 3: C6 Lane 4: C6 was treated with Pervanadate (10mM) for 1 hour Predicted band size: 149kDa Observed band size: 149kDa



Human appendix was stained with anti-PLCγ1 (Phospho Tyr783) (PT0758R) Rabbit antibody

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
PLC γ1 (Phospho Tyr783) (PT0758R) PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)