

BRD4 (PT1026R) PT™ Rabbit mAb

CatalogNo: YM8815 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, IP, ELISA

MW

- 152kD (Calculated)
152kD (Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

IHC 1:100-1:200

WB 1:2000-1:10000

IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

Basic Information

Clonality Monoclonal

Clone Number PT1026R

Immunogen Information

Immunogen The specific immunogen used to produce this antibody is proprietary information.

Specificity Endogenous

| Target Information

Gene name BRD4 HUNK1

Protein Name Bromodomain-containing protein 4 (Protein HUNK1)


Organism	Gene ID	UniProt ID
Human	23476 ;	O60885 ;
Mouse	57261 ;	Q9ESU6 ;

Cellular Localization Nucleus . Chromosome . Associates with acetylated chromatin (PubMed:21890894 , PubMed:16109376) . Released from chromatin upon deacetylation of histones that can be triggered by different signals such as activation of the JNK pathway or nocodazole treatment (PubMed:21890894 , PubMed:16109376) . Preferentially localizes to mitotic chromosomes , while it does not localizes to meiotic chromosomes (PubMed:21890894 , PubMed:16109376) . . ; [Isoform B]: Chromosome .


Tissue specificity Ubiquitously expressed.

Function Chromatin reader protein that recognizes and binds acetylated histones and plays a key role in transmission of epigenetic memory across cell divisions and transcription regulation. Remains associated with acetylated chromatin throughout the entire cell cycle and provides epigenetic memory for postmitotic G1 gene transcription by preserving acetylated chromatin status and maintaining high-order chromatin structure . During interphase , plays a key role in regulating the transcription of signal-inducible genes by associating with the P-TEFb complex and recruiting it to promoters. Also recruits P-TEFb complex to distal enhancers , so called anti-pause enhancers in collaboration with JMJD6. BRD4 and JMJD6 are required to form the transcriptionally active P-TEFb complex by displacing negative regulators such as HEXIM1 and 7SKsnRNA complex from P-TEFb , thereby transforming it into an active form that can then phosphorylate the C-terminal domain (CTD) of RNA polymerase II . Promotes phosphorylation of 'Ser-2' of the C-terminal domain (CTD) of RNA polymerase II . According to a report , directly acts as an atypical protein kinase and mediates phosphorylation of 'Ser-2' of the C-terminal domain (CTD) of RNA polymerase II; these data however need additional evidences in vivo . In addition to acetylated histones , also recognizes and binds acetylated RELA , leading to further recruitment of the P-TEFb complex and subsequent activation of NF-kappa-B . Also acts as a regulator of p53/TP53-mediated transcription: following phosphorylation by CK2 , recruited to p53/TP53 specific target promoters . ; [Isoform B]: Acts as a chromatin insulator in the DNA damage response pathway. Inhibits DNA damage response signaling by recruiting the condensin-2 complex to acetylated histones , leading to chromatin structure remodeling , insulating the region from DNA damage response by limiting spreading of histone H2AX/H2A.x phosphorylation.

| Validation Data



Rat kidney was stained with anti-BRD4 (PT1026R) Rabbit antibody



Various whole cell lysates were separated by 4-8% SDS-PAGE, and the membrane was blotted with anti-BRD4 (PT1026R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: HEK293 Lane 2: NIH-3T3 Lane 3: PC-12 Predicted band size: 152kDa Observed band size: 152kDa

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
BRD4 (PT1026R)
PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)