

# PLC γ2 (PT1152R) PT™ Rabbit mAb

CatalogNo: YM8884 Recombinant R

### **Key Features**

**Host Species** 

Rabbit

MW
• 148kD (Calculated)

148kD (Observed)

Reactivity

· Human, Mouse, Rat

Isotype

IgG,Kappa

**Applications** 

• WB,IHC,IF,IP,ELISA

### Recommended Dilution Ratios

IHC 1:200-1:1000 WB 1:2000-1:10000 IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

### **Storage**

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

### **Basic Information**

**Clonality** Monoclonal

Clone Number PT1152R

## Immunogen Information

**Specificity** Endogenous

# | Target Information

#### Gene name

PLCG2

#### **Protein Name**

1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-2

Organism	Gene ID	UniProt ID
Human	<u>5336</u> ;	<u>P16885;</u>
Mouse	<u>234779;</u>	Q8CIH5;
Rat	<u>29337;</u>	<u>P24135;</u>

#### Cellular Localization


intracellular,cytosol,plasma membrane,extracellular exosome,

**Tissue specificity** Lymph,Lymphoblast,Spleen,T-cell,

#### **Function**

Catalytic activity:1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate + H(2)O = 1D-myoinositol 1,4,5-trisphosphate + diacylglycerol.,cofactor:Calcium.,Function:The production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes. It is a crucial enzyme in transmembrane signaling., PTM: Phosphorylated on tyrosine residues; upon ligand-induced activation of a variety of growth factor receptors and immune system receptors. Increases phospholipase activity, similarity: Contains 1 C2 domain., similarity: Contains 1 PH domain., similarity: Contains 1 PI-PLC X-box domain., similarity: Contains 1 PI-PLC Y-box domain., similarity: Contains 1 SH3 domain., similarity: Contains 2 SH2 domains.,

## **Validation Data**



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-PLC y2 (PT1152R) antibody. The HRPconjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: A20 Predicted band size: 148kDa Observed band size: 148kDa

# | Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: PLC γ2 (PT1152R) PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents