

4E-BP1 (Phospho Thr70) (PT1203R) PT™ Rabbit mAb

CatalogNo: YM9045 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 13kD (Calculated)
15-20kD (Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

IHC 1:200-1:1000

WB 1:500-1:2000

IF 1:200-1:1000

ELISA 1:5000-1:20000

Basic Information

Clonality Monoclonal

Clone Number PT1203R

Immunogen Information

Immunogen The specific immunogen used to produce this antibody is proprietary information.

Specificity 4E-BP1 (Phospho Thr70) Antibody detects endogenous levels of 4E-BP1 protein only when phosphorylated at T70. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):TKtPP

Target Information

Gene name

EIF4EBP1

Protein Name

Eukaryotic translation initiation factor 4E-binding protein 1

Organism	Gene ID	UniProt ID
Human	1978;	Q13541;
Mouse	13685;	Q60876;
Rat	116636;	Q62622;

Cellular Localization

Nucleus


Tissue specificity

Colon ,Epithelium ,Lung ,Placenta ,Platelet ,

Function

Function:Regulates eIF4E activity by preventing its assembly into the eIF4F complex. Mediates the regulation of protein translation by hormones , growth factors and other stimuli that signal through the MAP kinase pathway. ,PTM:Phosphorylated on serine and threonine residues in response to insulin , EGF and PDGF. Phosphorylated upon DNA damage , probably by ATM or ATR. ,similarity:Belongs to the eIF4E-binding protein family. ,subunit:Nonphosphorylated EIF4EBP1 competes with EIF4G1/EIF4G3 to interact with EIF4E; insulin stimulated MAP-kinase (MAPK1 and MAPK3) phosphorylation of EIF4EBP1 causes dissociation of the complex allowing EIF4G1/EIF4G3 to bind and consequent initiation of translation. Rapamycin can attenuate insulin stimulation , mediated by FKBP. ,

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-E-BP1 (Phospho Thr70) (PT1203R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: C2C12 Lane 2: C2C12 was treated with Insulin (100 nM) for 15 minutes Predicted band size: 13kDa Observed band size: 15-20kDa

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:

**4E-BP1 (Phospho
Thr70) (PT1203R)
PT™ Rabbit mAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)