

HER2 (Phospho Tyr1248) /HER4 (Phospho Tyr1284) (PT1236R) PT™ Rabbit mAb

CatalogNo: YM9078 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IF, IP, ELISA

MW

- 147kD (Calculated)
180kD (Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

WB 1:2000-1:10000

IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

Basic Information

Clonality Monoclonal

Clone Number PT1236R

Immunogen Information

Immunogen The specific immunogen used to produce this antibody is proprietary information.

Specificity

HER2 (Phospho Tyr1248) /HER4 (Phospho Tyr1284) Antibody detects endogenous levels of HER2/HER4 protein only when phosphorylated at Y1248(HER2)/Tyr1284(HER4).The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):PEyLG

Target Information


Gene name ERBB2 HER2 ERBB4 HER4

Protein Name Receptor tyrosine-protein kinase erbB-2/Receptor tyrosine-protein kinase erbB-4

Organism	Gene ID	UniProt ID
Human	2064 ; 2066 ;	P04626 ; Q15303 ;
Mouse	13866 ; 13869 ;	P70424 ; Q61527 ;
Rat	59323 ;	P06494 ; Q62956 ;

Function Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. ,Disease:Defects in ERBB2 are associated with familial glioma of brain [MIM:137800]; also called glioblastoma multiforme. Gliomas are central nervous system neoplasms derived from glial cells and comprise astrocytomas , glioblastoma multiforme , oligodendrogliomas , and ependymomas. ,Disease:Defects in ERBB2 are associated with gastric cancer [MIM:137215]; also known as hereditary familial diffuse gastric cancer (HDGC) . ,Disease:Defects in ERBB2 are associated with lung cancer [MIM:211980]; also called adenocarcinoma of lung. ,Disease:Defects in ERBB2 are associated with ovarian cancer [MIM:167000]. Ovarian cancer is the leading cause of death from gynecologic malignancy. It is characterized by advanced presentation with loco-regional dissemination in the peritoneal cavity and the rare incidence of visceral metastases. These typical features relate to the biology of the disease , which is a principal determinant of outcome. ,Function:Essential component of a neuregulin-receptor complex , although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Not activated by EGF , TGF-alpha and amphiregulin. ,online information:ERBB2 entry ,polymorphism:There are four alleles due to the variations in positions 654 and 655. Allele B1 (Ile-654/Ile-655) has a frequency of 0.782; allele B2 (Ile-654/Val-655) has a frequency of 0.206; allele B3 (Val-654/Val-655) has a frequency of 0.012. ,PTM:Ligand-binding increases phosphorylation on tyrosine residues. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. EGF receptor subfamily. ,similarity:Contains 1 protein kinase domain. ,subunit:Heterodimer with each of the other ERBB receptors (Potential) . Interacts with PRKCABP and PLXNB1. Part of a complex with EGFR and either PIK3C2A or PIK3C2B. May interact with PIK3C2B when phosphorylated on Tyr-1196. Interacts with MEMO when phosphorylated on Tyr-1248. Interacts with MUC1. Stimulation by heregulin (HRG) in breast cancer cell lines induces binding of MUC1 with gamma-catenin. ,

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-HER2 (Phospho Tyr1248) /HER4 (Phospho Tyr1284) (PT1236R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: HeLa Lane 2: HeLa serum was starved for 4 hours and treated with EGF (200ng/ml) for 15 minutes Predicted band size: 147kDa Observed band size: 180kDa

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
HER2 (Phospho Tyr1248) /HER4 (Phospho Tyr1284) (PT1236R) PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)