

NDRG1 (Phospho Thr346) (PT1585R) PT™ Rabbit mAb

CatalogNo: YM9427 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 43kD (Calculated)
50kD (Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

IHC 1:200-1:1000

WB 1:2000-1:10000

IF 1:200-1:1000

ELISA 1:5000-1:20000

Basic Information

Clonality Monoclonal

Clone Number PT1585R

Immunogen Information

Immunogen The specific immunogen used to produce this antibody is proprietary information.

Specificity NDRG1 (Phospho Thr346) antibody detects endogenous levels of Human Mouse Rat NDRG1 (phospho-Thr346) .The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):SHtSE

| **Target Information**

Gene name NDRG1 CAP43 DRG1 RTP

Protein Name NDRG1 (Thr346)

Organism	Gene ID	UniProt ID
Human	10397;	Q92597;
Mouse	17988;	Q62433;
Rat	299923;	Q6JE36;


Cellular Localization Cytoplasm , cytosol. Cytoplasm , cytoskeleton , microtubule organizing center , centrosome. Nucleus. Cell membrane. Mainly cytoplasmic but differentially localized to other regions. Associates with the plasma membrane in intestinal epithelia and lactating mammary gland. Translocated to the nucleus in a p53/TP53-dependent manner. In prostate epithelium and placental chorion , located in both the cytoplasm and in the nucleus. No nuclear localization in colon epithelium cells. In intestinal mucosa , prostate and renal cortex , located predominantly adjacent to adherens junctions. Cytoplasmic with granular staining in proximal tubular cells of the kidney and salivary gland ducts. Recruits to the membrane of recycling/sorting and late endosomes via binding to phosphatidylinositol 4-phosphate. Associates with microtubules. Colocalizes with TUBG1 in the centrosome. Cytoplasmic location increased with hypoxia. Phosphorylated form found associated with centromeres during S-phase of mitosis and with the plasma membrane.

Tissue specificity Ubiquitous; expressed most prominently in placental membranes and prostate , kidney , small intestine , and ovary tissues. Also expressed in heart , brain , skeletal muscle , lung , liver and pancreas. Low levels in peripheral blood leukocytes and in tissues of the immune system. Expressed mainly in epithelial cells. Also found in Schwann cells of peripheral neurons. Reduced expression in adenocarcinomas compared to normal tissues. In colon , prostate and placental membranes , the cells that border the lumen show the highest expression.


Function

Disease:Defects in NDRG1 are the cause of Charcot-Marie-Tooth disease type 4D (CMT4D) [MIM:601455]; also known as hereditary motor and sensory neuropathy Lom type (HMSNL) . CMT4D is a recessive form of Charcot-Marie-Tooth disease , the most common inherited disorder of the peripheral nervous system. Charcot-Marie-Tooth disease is classified in two main groups on the basis of electrophysiologic properties and histopathology: primary peripheral demyelinating neuropathy and primary peripheral axonal neuropathy. Demyelinating CMT neuropathies are characterized by severely reduced nerve conduction velocities (less than 38 m/sec) , segmental demyelination and remyelination with onion bulb formations on nerve biopsy , slowly progressive distal muscle atrophy and weakness , absent deep tendon reflexes , and hollow feet. By convention , autosomal recessive forms of demyelinating Charcot-Marie-Tooth disease are designated CMT4. ,Function:May have a growth inhibitory role. ,induction:By homocysteine , 2-mercaptoethanol , tunicamycin in endothelial cells. Induced approximately 20-fold during in vitro differentiation of the colon carcinoma cell lines HT29-D4 and Caco-2. Induced by nickel compounds in all tested cell lines. The primary signal for its induction is an elevation of free intracellular calcium ion caused by nickel ion exposure. Okadaic acid , a serine/threonine phosphatase inhibitor , induced its expression more rapidly and more efficiently than nickel. ,similarity:Belongs to the NDRG family. ,subcellular location:Whereas in prostate epithelium and placental chorion it is located in both the cytoplasm and the nucleus , nuclear staining is not observed in colon epithelium cells. Instead its localization changes from the cytoplasm to the plasma membrane during differentiation of colon carcinoma cell lines in vitro. ,tissue specificity:Ubiquitous; expressed most prominently in placental membranes and prostate , kidney , small intestine , and ovary tissues. Reduced expression in adenocarcinomas compared to normal tissues. In colon , prostate and placental membranes , the cells that border the lumen show the highest expression. ,


Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-NDRG1 (Phospho Thr346) (PT1585R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: HT-1080 Lane 2: PC-3 Lane 3: RAW264.7 Lane 4: Rat brain Predicted band size: 43kDa Observed band size: 50kDa



Human tonsil was stained with anti-NDRG1 (Phospho Thr346) (PT1585R) Rabbit antibody



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-NDRG1 (Phospho Thr346) (PT1585R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: Jurkat Lane 2: Jurkat was treated with Calyculin A (0.1 μ M) for 45 minutes Lane 3: C6 Lane 4: C6 was treated with Calyculin A (100ng/mL) for 30 minutes Predicted band size: 43kDa Observed band size: 50kDa

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
NDRG1 (Phospho Thr346) (PT1585R) PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)