

β Catenin (Phospho Ser552) (PT1629R) PT™ Rabbit mAb

CatalogNo: YM9471 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, IP, ELISA

MW

- 85kD (Calculated)
100kD (Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

IHC 1:200-1:1000

WB 1:2000-1:10000

IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

Basic Information

Clonality Monoclonal

Clone Number PT1629R

Immunogen Information

Immunogen The specific immunogen used to produce this antibody is proprietary information.


Specificity

β Catenin (Phospho Ser552) Antibody detects endogenous levels of Catenin-β protein only when phosphorylated at S552.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RTsMG


Target Information

Gene name	CTNNB1 CTNNB OK/SW-cl.35 PRO2286		
Protein Name	Catenin-β;b-catenin;Beta catenin;Beta-catenin;Cadherin associated protein;Catenin (cadherin associated protein) , beta 1, 88 kDa;Catenin beta 1;Catenin beta-1;CATNB;CHBCAT;CTNB1_HUMAN;CTNNB;CTNNB1;DKFZp686D0253;FLJ25606;FLJ37923;OTTHUMP00000162082;OTTHUMP00000165222;OTTHUMP00000165223;OTTHUMP00000209288;OTTHUMP00000209289		
	Organism	Gene ID	UniProt ID
	Human	1499;	P35222;
	Mouse	12387;	Q02248;
	Rat	84353;	Q9WU82;
Cellular Localization	Cytoplasm . Nucleus . Cytoplasm , cytoskeleton . Cell junction , adherens junction . Cell junction . Cell membrane . Cytoplasm , cytoskeleton , microtubule organizing center , centrosome. Cytoplasm , cytoskeleton , spindle pole . Cell junction , synapse . Cytoplasm , cytoskeleton , cilium basal body . Colocalized with RAPGEF2 and TJP1 at cell-cell contacts (By similarity) . Cytoplasmic when it is unstabilized (high level of phosphorylation) or bound to CDH1. Translocates to the nucleus when it is stabilized (low level of phosphorylation) . Interaction with GLIS2 and MUC1 promotes nuclear translocation. Interaction with EMD inhibits nuclear localization. The majority of beta-catenin is localized to the cell membrane. In interphase , colocalizes with CROCC between CEP250 puncta at the proximal end of centrioles , and this localization is dependent on CROCC and CEP250. In mitosis , when NEK2 activity increases , it localizes to centrosomes at spindle poles independent of CROCC. Colocalizes with CDK5 in the cell-cell contacts and plasma membrane of undifferentiated and differentiated neuroblastoma cells. Interaction with FAM53B promotes translocation to the nucleus (PubMed:25183871) . .		
Tissue specificity	Expressed in several hair follicle cell types: basal and peripheral matrix cells , and cells of the outer and inner root sheaths. Expressed in colon. Present in cortical neurons (at protein level) . Expressed in breast cancer tissues (at protein level) (PubMed:29367600) .		
Function	Disease:A chromosomal rearrangement involving CTNNB1 may be a cause of salivary gland pleiomorphic adenomas (PA) [181030]. Pleiomorphic adenomas are the most common benign epithelial tumors of the salivary gland. Translocation t (3;8) (p21;q12) with PLAG1 .,Disease:Activating mutations in CTNNB1 have oncogenic activity resulting in tumor development. Somatic mutations are found in various tumor types , including colon cancers , ovarian and prostate carcinomas , hepatoblastoma (HB) , hepatocellular carcinoma (HCC) . HBs are malignant embryonal tumors mainly affecting young children in the first three years of life. .,Disease:Defects in CTNNB1 are a cause of medulloblastoma (MDB) [MIM:155255]. MDB is a malignant , invasive embryonal tumor of the cerebellum with a preferential manifestation in children. .,Disease:Defects in CTNNB1 are a cause of pilomatixoma (PTR) [MIM:132600]; a common benign skin tumor. .,Disease:Defects in CTNNB1 are associated with colorectal cancer (CRC) [MIM:114500]. .,Disease:Defects in CTNNB1 are associated with ovarian cancer [MIM:167000]. Ovarian cancer is the leading cause of death from gynecologic malignancy. It is characterized by advanced presentation with loco-regional dissemination in the peritoneal cavity and the rare incidence of visceral metastases. These typical features relate to the biology of the disease , which is a principal determinant of outcome. .,Function:Involved in the regulation of cell adhesion and in signal transduction through the Wnt pathway. .,online information:Beta-catenin entry ,PTM:EGF stimulates tyrosine phosphorylation. Phosphorylation on Tyr-654 decreases CDH1 binding and enhances TBP binding. .,PTM:Phosphorylation by GSK3B requires prior phosphorylation of Ser-45 by another kinase. Phosphorylation proceeds then from Thr-41 to Ser-37 and Ser-33. .,PTM:Ubiquitinated by a E3 ubiquitin ligase complex containing UBE2D1 , SIAH1 , CACYBP/SIP , SKP1A , APC and TBL1X (Probable) . Its ubiquitination leads to its subsequent proteasomal degradation. .,similarity:Belongs to the beta-catenin family. .,similarity:Contains 12 ARM repeats. .,subcellular location:Cytoplasmic when it is unstabilized (high level of phosphorylation) or bound to CDH1. Translocates to the nucleus when it is stabilized (low level of phosphorylation) . Interaction with GLIS2 and MUC1 promotes nuclear translocation. .,subunit:Two separate pools are found in the cytoplasm: one is PSEN1/cadherin/catenin complex which anchors to the actin cytoskeleton. The other pool is part of a large complex containing AXIN1 , AXIN2 , APC , CSNK1A1 and GSK3B that promotes phosphorylation on N-terminal Ser and Thr residues and ubiquitination of CTNNB1 via BTRC and its subsequent degradation by the proteasome. Wnt-dependent activation of DVL antagonizes the action of GSK3B. When GSK3B activity is inhibited the complex dissociates , CTNNB1 is dephosphorylated and is no longer targeted for destruction. The stabilized protein translocates to the nucleus , where it binds TCF/LEF-1 family members , TBP , BCL9 and possibly also RUVBL1 and CHD8. Binds CTNNBIP and EP300. CTNNB1 forms a ternary complex with LEF1 and EP300 that is disrupted by CTNNBIP1 binding (By similarity) . Interacts with TAX1BP3 (via the PDZ domain) ; this interaction inhibits the transcriptional activity of CTNNB1 (By similarity) . Interacts with AJAP1 , BAIAP1 , CARM1 , CTNNA3 , CXADR and PCDH11Y. Binds SLC9A3R1. Interacts with GLIS2 and MUC1. Interacts with SLC30A9. Interacts with XIRP1 (By similarity) . Interacts with PTPRU (via the cytoplasmic juxtamembrane domain) . .,tissue specificity:Expressed in several hair follicle cell types: basal and peripheral matrix cells , and cells of the outer and inner root sheaths. Expressed in colon. ,		

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-β Catenin (Phospho Ser552) (PT1629R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: NIH-3T3 Lane 2: NIH-3T3 was treated with Forskolin (30uM) for 1 hour Lane 3: PC-12 Lane 4: PC-12 was treated with Forskolin (30uM) for 1 hour Predicted band size: 85kDa Observed band size: 100kDa



Human liver was stained with anti-β Catenin (Phospho Ser552) (PT1629R) Rabbit antibody

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
β Catenin (Phospho Ser552) (PT1629R) PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)