

## SMG9 Rabbit pAb

CatalogNo: YN0146

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, ELISA

#### MW

- 57kD (Observed)

#### Isotype

- IgG

### Recommended Dilution Ratios

**WB 1:500-2000**

**ELISA 1:5000-20000**

### Storage

#### Storage\*

-15°C to -25°C/1 year (Do not lower than -25°C)

#### Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Basic Information

#### Clonality

Polyclonal

### Immunogen Information

#### Immunogen

Synthesized peptide derived from human protein . at AA range: 70-150

#### Specificity

SMG9 Polyclonal Antibody detects endogenous levels of protein.


### Target Information

#### Gene name

SMG9 C19orf61

<b>Protein Name</b>	Protein SMG9 (Protein smg-9 homolog)		
	<b>Organism</b>	<b>Gene ID</b>	<b>UniProt ID</b>
	Human	<a href="#">56006</a> ;	<a href="#">Q9H0W8</a> ;
	Mouse		<a href="#">Q9DB90</a> ;
	Rat		<a href="#">Q5PQS6</a> ;
<b>Cellular Localization</b>	intracellular,cytosol,		
<b>Tissue specificity</b>	Brain,Epithelium,Muscle,Teratocarcinoma,		
<b>Function</b>	<p>Function:Component of the SMG1C complex, a mRNA surveillance complex that recognizes and degrades mRNAs containing premature translation termination codons (PTCs) via the nonsense-mediated mRNA decay (NMD). The complex probably acts by associating with ribosomes during tranlation termination on mRNPs. If an exon junction complex (EJC) is located 50-55 or more nucleotides downstream from the termination codon, smg1 phosphorylates upf1/rent1, triggering nonsense-mediated mRNA decay (NMD). In the SMG1C complex, it is required for the efficient association between smg1 and smg8.,PTM:Phosphorylated by SMG1.,similarity:Belongs to the SMG9 family.,subunit:Component of the SMG1C complex, at least composed of SMG1, SMG8 and SMG9. The SMG1C complex is then recruited on premature translation termination codons (PTCs) to form the ribosome:SURF complex, at least composed of ERF1, ERF3 (ERF3A or ERF3B), EEF2, UPF1/RENT1, SMG1, SMG8 and SMG9.,</p>		

Validation Data



Western Blot analysis of HEK293 lysis, using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000

## | Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**SMG9 Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)