

# KIF4B Rabbit pAb

CatalogNo: YN0596

### **Key Features**

**Host Species** 

Rabbit

Reactivity

Human

**Applications** WB,ELISA

MW

135kD (Observed)

Isotype IgG

#### Recommended Dilution Ratios

WB 1:500-2000

ELISA 1:5000-20000

## Storage

Storage\*

-15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** 

Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

### **I** Basic Information

Clonality

Polyclonal

## Immunogen Information

**Immunogen** 

Synthesized peptide derived from part region of human protein

**Specificity** 

KIF4B Polyclonal Antibody detects endogenous levels of protein.

## **| Target Information**

**Gene name** 

KIF4B

#### **Protein Name**

Chromosome-associated kinesin KIF4B (Chromokinesin-B)

Organism	Gene ID	UniProt ID
Human	<u>285643</u> ;	Q2VIQ3;

#### Cellular Localization


Nucleus matrix . Cytoplasm, cytoskeleton .

**Tissue specificity** Specifically expressed in testis.

#### **Function**

Function:Motor protein that translocates PRC1 to the plus ends of interdigitating spindle microtubules during the metaphase to anaphase transition, an essential step for the formation of an organized central spindle midzone and midbody and for successful cytokinesis. May play a role in mitotic chromosomal positioning and bipolar spindle stabilization.,sequence Caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the kinesin-like protein family.,similarity:Belongs to the kinesin-like protein family. Chromokinesin subfamily.,similarity:Contains 1 kinesin-motor domain.,subcellular location:Not present in the nucleolus. In early mitosis, associated with the mitotic spindle, in anaphase, localized to the spindle midzone and, in telophase and cytokinesis, to the midbody. In late cytokinesis, found in the center of the midbody. Associated with chromosomes at all stages of mitosis.,subunit:Interacts with unphosphorylated PRC1 during late mitosis.,tissue specificity:Highly expressed in hematopoietic tissues, fetal liver, spleen, thymus and adult thymus and bone marrow. Lower levels are found in heart, testis, kidney, colon and lung.,tissue specificity:Specifically expressed in testis.,

### **Validation Data**



Western Blot analysis of HEK293 lysis, using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000

## | Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **KIF4B Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents