

DHB12 Rabbit pAb

CatalogNo: YN0706

Key Features

Host Species

Rabbit

Reactivity

· Human, Mouse, Rat

ApplicationsWB,ELISA

MW

34kD (Observed)

IsotypeIgG

Recommended Dilution Ratios

WB 1:500-2000 ELISA 1:5000-20000

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from part region of human protein

Specificity DHB12 Polyclonal Antibody detects endogenous levels of protein.

| Target Information

Gene name HSD17B12

Protein Name

Estradiol 17-beta-dehydrogenase 12 (17-beta-hydroxysteroid dehydrogenase 12) (17-beta-HSD 12) (3-ketoacyl-CoA reductase) (KAR) (EC 1.3.1.-)

Organism	Gene ID	UniProt ID	
Human	<u>51144;</u>	<u>Q53GQ0</u> ;	
Mouse		<u>070503;</u>	
Rat		<u>Q6P7R8;</u>	

Cellular Localization


Endoplasmic reticulum membrane; Multi-pass membrane protein.

Tissue specificity Expressed in most tissues tested. Highly expressed in the ovary and mammary. Expressed in platelets.

Function

Catalytic activity: Estradiol-17-beta + NAD(P)(+) = estrone + NAD(P)H., Domain: The di-lysine motif confers endoplasmic reticulum localization for type I membrane proteins., Function: Catalyzes the transformation of estrone (E1) into estradiol (E2), suggesting a central role in estrogen formation. Its strong expression in ovary and mammary gland suggest that it may constitute the major enzyme responsible for the conversion of E1 to E2 in women. Also has 3-ketoacyl-CoA reductase activity, reducing both long chain 3-ketoacyl-CoAs and long chain fatty acyl-CoAs, suggesting a role in long fatty acid elongation.,pathway:Lipid metabolism; fatty acid biosynthesis.,pathway:Steroid biosynthesis; estrogen biosynthesis., similarity: Belongs to the short-chain dehydrogenases/reductases (SDR) family. 17-beta-HSD 3 subfamily., tissue specificity: Expressed in most tissues tested. Highly expressed in the ovary and mammary. Expressed in platelets.,

I Validation Data



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night

I Contact information

Orders: order.cn@immunoway.com support.cn@immunoway.com Support:

400-8787-807(China) Telephone:

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **DHB12** Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents