

MUC7 Rabbit pAb

CatalogNo: YN0882

Key Features

Host Species

- Rabbit

Reactivity

- Human

Applications

- WB,ELISA

MW

- 41kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-2000**ELISA 1:5000-20000**

Storage

Storage*

-15°C to -25°C/1 year(Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality

Polyclonal

Immunogen Information

Immunogen

Synthesized peptide derived from human protein . at AA range: 40-120

Specificity

MUC7 Polyclonal Antibody detects endogenous levels of protein.


Target Information

Gene name

MUC7 MG2

Protein Name	Mucin-7 (MUC-7) (Apo-MG2) (Salivary mucin-7)		
	Organism	Gene ID	UniProt ID
	Human	4589;	Q8TAX7;
Cellular Localization	Secreted .		
Tissue specificity	Expressed in salivary gland tissues and only in those that contain mucous acinar cells (e.g. sublingual and submandibular glands) and not in salivary glands containing only serous acinar cells (e.g. parotid gland).		
Function	<p>Disease:MUC7 alleles are associated with susceptibility to asthma [MIM:600807]. The MUC7*5 allele is rarer in the atopic asthmatics than in the atopic non-asthmatics. Comparison of all atopic individuals with all nonatopic shows no difference, while comparison of all asthmatic individuals with all non-asthmatic shows that the asthmatic group has reduced MUC7*5 frequency. The significantly lower frequency of the MUC7*5 allele in individuals with atopic asthma was explained by the possible association between alleles and different interactions with bacteria, since the glycosylated domain is thought to be responsible, at least in part, for the bacterial binding that allows bacteria to be cleared from the epithelial surfaces.,Function:May function in a protective capacity by promoting the clearance of bacteria in the oral cavity and aiding in mastication, speech, and swallowing. Binds P.aeruginosa pili.,polymorphism:The most common allele, MUC7*6, contains a tandem repeat domain comprising 6 repeats (shown here) each composed of 23 amino acids. These repeats are very similar but not identical. In a large cohort of 375 individuals from a variety of ethnic backgrounds, three different alleles were detected, MUC7*6 being the most common, in all populations studied, followed by MUC7*5 (5 repeats), with frequency varying from 0.05 in Africans to 0.22 in East Asians. MUC7*8 (8 repeats), a novel rare allele, was identified in 1 Northern European individual.,PTM:N- and O-glycosylated. Contains fucose, mannose, galactose, N-acetylglucosamine and N-acetylgalactosamine.,subunit:Monomer.,tissue specificity:Expressed in salivary gland tissues and only in those that contain mucous acinar cells (e.g. sublingual and submandibular glands) and not in salivary glands containing only serous acinar cells (e.g. parotid gland).,</p>		

| Validation Data



Western blot analysis of lysates from PC12 cells, primary antibody was diluted at 1:1000, 4°over night

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
MUC7 Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)