

#### www.immunoway.com.cn

# MYH2 Rabbit pAb

CatalogNo: YN0898

# Key Features

Host Species • Rabbit	Reactivity • Human	Applications <ul> <li>WB,ELISA</li> </ul>
MW • 213kD (Observed)	lsotype • lgG	

#### **Recommended Dilution Ratios**

WB 1:500-2000 ELISA 1:5000-20000

#### **Storage**

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

#### **Basic Information**

Clonality Polyclonal

## Immunogen Information

ImmunogenSynthesized peptide derived from human protein . at AA range: 760-840SpecificityMYH2 Polyclonal Antibody detects endogenous levels of protein.

## **Target Information**

Gene name MYH2 MYHSA2

Protein Name	Myosin-2 (Myosin heavy chain 2) (Myosin heavy chain 2a) (MyHC-2a) (Myosin heavy chain
	lla) (MyHC-lla) (Myosin heavy chain, skeletal muscle, adult 2)

Organism	Gene ID	UniProt ID
Human	<u>4620;</u>	<u>Q9UKX2;</u>

**Cellular** Cytoplasm, myofibril. Thick filaments of the myofibrils.

Tissue specificity Cerebellum, Muscle pool- 2 tissues- cardiac and skeletal muscle., Skeletal muscle,

**Function** Disease:Defects in MYH2 are the cause of inclusion body myopathy type 3 (IBM3) [MIM:605637]. Hereditary inclusion body myopathies constitute a group of neuromuscular disorders characterized by slowly progressive distal and proximal weakness and a typical muscle pathology including rimmed vacuoles and filamentous inclusions. IBM3 is a variant of hereditary inclusion body myopathies and is characterized by autosomal dominant myopathy with joint contracture, ophthalmoplegia and rimmed vacuoles. Morphological analysis of muscle biopsies from patients indicate that the type 2A fibers frequently were abnormal, whereas other fiber types appeared normal.,Domain:The rodlike tail sequence is highly repetitive, showing cycles of a 28-residue repeat pattern composed of 4 heptapeptides, characteristic for alpha-helical coiled coils., Function: Muscle contraction. Required for cytoskeleton organization., miscellaneous: Each myosin heavy chain can be split into 1 light meromyosin (LMM) and 1 heavy meromyosin (HMM). It can later be split further into 2 globular subfragments (S1) and 1 rod-shaped subfragment (S2)., similarity: Contains 1 IQ domain., similarity: Contains 1 myosin head-like domain., subcellular location: Thick filaments of the myofibrils., subunit: Muscle myosin is a hexameric protein that consists of 2 heavy chain subunits (MHC), 2 alkali light chain subunits (MLC) and 2 regulatory light chain subunits (MLC-2).,

#### Validation Data

Localization

#### **Contact information**

Orders:	order.cn@immunoway.com
Support:	support.cn@immunoway.com
Telephone:	400-8787-807(China)
Website:	http://www.immunoway.com.cn
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **MYH2 Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents