

2A5G Rabbit pAb

CatalogNo: YN1279

Key Features

Host Species Reactivity Applications
• Rabbit • Human, Mouse • WB, ELISA

MW Isotype • 57kD (Observed) • IgG

Recommended Dilution Ratios

WB 1:500-2000 ELISA 1:5000-20000

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human protein . at AA range: 360-440

Specificity 2A5G Polyclonal Antibody detects endogenous levels of protein.

| Target Information

Gene name PPP2R5C KIAA0044

Protein Name

Serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit gamma isoform (PP2A B subunit isoform B'-gamma) (PP2A B subunit isoform B56-gamma) (PP2A B subunit isoform PR61-gamma) (PP2A B subunit isoform R5-gamma) (Renal carcinoma antigen NY-REN-29)

Organism	Gene ID	UniProt ID	
Human	<u>5527;</u>	<u>Q13362</u> ;	
Mouse		<u>Q60996</u> ;	

Cellular Localization

Nucleus. Chromosome, centromere.

Tissue specificity Highest levels in heart, skeletal muscle and brain. Lower levels in pancreas, kidney, lung and placenta. Very low levels in liver.

Function

Function: The B regulatory subunit might modulate substrate selectivity and catalytic activity, and also might direct the localization of the catalytic enzyme to a particular subcellular compartment.,PTM:Isoform Gamma-3 is phosphorylated on serine residues while isoform Gamma-1 is not., similarity: Belongs to the phosphatase 2A regulatory subunit B56 family,, subunit: PP2A consists of a common heterodimeric core enzyme, composed of a 36 kDa catalytic subunit (subunit C) and a 65 kDa constant regulatory subunit (PR65 or subunit A), that associates with a variety of regulatory subunits. Proteins that associate with the core dimer include three families of regulatory subunits B (the R2/B/PR55/B55, R3/B"/PR72/PR130/PR59 and R5/B'/B56 families), the 48 kDa variable regulatory subunit, viral proteins, and cell signaling molecules. Interacts with SGOL1., tissue specificity: Highest levels in heart, skeletal muscle and brain. Lower levels in pancreas, kidney, lung and

placenta. Very low levels in liver.,

Validation Data

Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: 2A5G Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents