

SPAG5 Rabbit pAb

CatalogNo: YN1358

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, ELISA

MW

- 131kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-2000**ELISA 1:5000-20000**

Storage

Storage*

-15°C to -25°C/1 year (Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality

Polyclonal

Immunogen Information

Immunogen

Synthesized peptide derived from part region of human protein

Specificity

SPAG5 Polyclonal Antibody detects endogenous levels of protein.

Target Information

Gene name

SPAG5

Protein Name	Sperm-associated antigen 5 (Astrin) (Deepest) (Mitotic spindle-associated protein p126) (MAP126)		
	Organism	Gene ID	UniProt ID
	Human	10615;	Q96R06;
	Mouse		Q7TME2;
Cellular Localization	Cytoplasm . Cytoplasm, cytoskeleton . Cytoplasm, cytoskeleton, spindle . Cytoplasm, cytoskeleton, spindle pole . Chromosome, centromere, kinetochore . Midbody. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasmic granule. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite . Colocalizes with PCM1 at centriolar satellites throughout the cell cycle (PubMed:26297806). In a punctate pattern in interphase cells. During mitosis, detected at spindle poles during prophase, throughout the spindle in metaphase and anaphase, and at midzone microtubules in anaphase and telophase (PubMed:27462074). Efficient targeting to the mitotic spindle may depend upon phosphorylation by GSK3B. Detected on kinetochores of chromosomes that have congressed. The astrin (SPAG5)-kinastrin (SKAP) complex localizes to the microtubule plus ends (By similarity). In non-mitotic non-stressed cells, shows a microtubuli pattern. In arsenite-stressed cells, accumulates in stress granules. .		
Tissue specificity	Highly expressed in testis. Detected at low levels in placenta, liver, pancreas, thymus and colon.		
Function	Function:Necessary for normal spindle formation during mitosis.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,subcellular location:In a punctate pattern in interphase cells. During mitosis, detected at spindle poles during prophase, throughout the spindle in metaphase and anaphase, and at midzone microtubules in anaphase and telophase. Detected on kinetochores of chromosomes that have congressed.,subunit:Homodimer, with a globular head domain and a long stalk. Homo-oligomer; the globular head domains associate, resulting in aster-like structures. Binds to microtubules in the mitotic spindle.,tissue specificity:Highly expressed in testis. Detected at low levels in placenta, liver, pancreas, thymus and colon.,		

| Validation Data

| Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
SPAG5 Rabbit pAb