

TNKS1 Rabbit pAb

CatalogNo: YN2269

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, ELISA

MW

- 145kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-2000**ELISA 1:5000-20000**

Storage

Storage*

-15°C to -25°C/1 year (Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality

Polyclonal

Immunogen Information

Immunogen

Synthesized peptide derived from part region of human protein

Specificity

TNKS1 Polyclonal Antibody detects endogenous levels of protein.

Target Information

Gene name

TNKS PARP5A PARPL TIN1 TINF1 TNKS1

Protein Name	Tankyrase-1 (TANK1) (ADP-ribosyltransferase diphtheria toxin-like 5) (ARTD5) (Poly [ADP-ribose] polymerase 5A) (TNKS-1) (TRF1-interacting ankyrin-related ADP-ribose polymerase) (Tankyrase I)		
	Organism	Gene ID	UniProt ID
	Human	8658 ;	O95271 ;
	Mouse		Q6PFX9 ;
Cellular Localization	Cytoplasm . Golgi apparatus membrane ; Peripheral membrane protein . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Nucleus, nuclear pore complex . Chromosome, telomere . Cytoplasm, cytoskeleton, spindle pole . Associated with the Golgi and with juxtanuclear SLC2A4/GLUT4-vesicles (PubMed:22864114). A minor proportion is also found at nuclear pore complexes and around the pericentriolar matrix of mitotic centromeres (PubMed:10523501). During interphase, a small fraction of TNKS is found in the nucleus, associated with TERF1 (PubMed:12768206). Localizes to spindle poles at mitosis onset via interaction with NUMA1 (PubMed:12080061). .		
Tissue specificity	Ubiquitous; highest levels in testis.		
Function	Catalytic activity:NAD(+) + (ADP-D-ribosyl)(n)-acceptor = nicotinamide + (ADP-D-ribosyl)(n+1)-acceptor.,Function:May regulate vesicle trafficking and modulate the subcellular distribution of SLC2A4/GLUT4-vesicles. Has PARP activity and can modify TERF1, and thereby contribute to the regulation of telomere length.,PTM:ADP-ribosylated (- auto).,PTM:Phosphorylated on serine residues by MAPK kinases upon insulin stimulation.,similarity:Contains 1 PARP catalytic domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,similarity:Contains 15 ANK repeats.,subcellular location:Associated with the Golgi and with juxtanuclear SLC2A4/GLUT4-vesicles. A minor proportion is also found at nuclear pore complexes and around the pericentriolar matrix of mitotic centromeres. During interphase, a small fraction of TNKS is found in the nucleus, associated with TERF1.,subunit:Oligomerizes and associates with TNKS2. Interacts with the cytoplasmic domain of LNPEP/Otase in SLC2A4/GLUT4-vesicles. Binds to the N-terminus of telomeric TERF1 via the ANK repeats. Found in a complex with POT1; TERF1 and TIN2.,tissue specificity:Ubiquitous; highest levels in testis.,		

Validation Data

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
TNKS1 Rabbit pAb

