

HMGA1 Rabbit pAb

CatalogNo: YN2897

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA

MW

- 11kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-2000

ELISA 1:5000-20000

Storage

Storage*

-15°C to -25°C/1 year (Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality

Polyclonal

Immunogen Information

Immunogen

Synthesized peptide derived from part region of human protein AA range: 1-50

Specificity

HMGA1 Polyclonal Antibody detects endogenous levels of protein.

Target Information

Gene name

HMGA1 HMG1Y

Protein Name	High mobility group protein HMG-I/HMG-Y (HMG-I(Y)) (High mobility group AT-hook protein 1) (High mobility group protein A1) (High mobility group protein R)		
	Organism	Gene ID	UniProt ID
	Human	3159;	P17096;
	Mouse	111241;	P17095;
	Rat	117062;	Q8K585;
Cellular Localization	Nucleus. Chromosome.		
Tissue specificity	B-cell,Colon,Epithelium,Kidney,Liver,Mammary gland,Muscle,Pancreas,Placenta		
Function	<p>Disease:A chromosomal aberration involving HMGA1 is found in pulmonary chondroid hamartoma. Translocation t(6;14)(p21;q23-24) with RAD51L1.,Function:HMG-I/Y bind preferentially to the minor groove of A+T rich regions in double stranded DNA. It is suggested that these proteins could function in nucleosome phasing and in the 3'-end processing of mRNA transcripts. They are also involved in the transcription regulation of genes containing, or in close proximity to A+T-rich regions.,mass spectrometry:With 1 acetyl and 2 phosphate groups PubMed:15302935,mass spectrometry:With 1 acetyl and 3 phosphate groups PubMed:15302935,mass spectrometry:With 1 acetyl, 1 methyl and 2 phosphate groups PubMed:15302935,mass spectrometry:With 1 acetyl, 1 methyl and 3 phosphate groups PubMed:15302935,mass spectrometry:With 1 acetyl, 2 methyl and 2 phosphate groups PubMed:15302935,mass spectrometry:With 1 acetyl, 2 methyl and 3 phosphate groups PubMed:15302935,PTM:Constitutively phosphorylated on two or three sites. Phosphorylated upon DNA damage, probably by ATM or ATR. Hyperphosphorylated at early stages of apoptosis, followed by dephosphorylation and methylation, which coincides with chromatin condensation. Isoform HMG-Y can be phosphorylated by HIPK2.,PTM:HMG-Y is not methylated.,PTM:Methylation at Arg-58 is mutually exclusive with methylation at Arg-60.,similarity:Belongs to the HMGA family.,similarity:Contains 3 A.T hook DNA-binding domains.,subunit:Interacts with HIPK2 (By similarity). Interacts with HIV-1 pre-integration complex.,</p>		

| Validation Data

| Contact information

Orders:

order.cn@immunoway.com

Support:

support.cn@immunoway.com

Telephone:

400-8787-807(China)

Website:

http://www.immunoway.com.cn

Address:

2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

HMGA1 Rabbit pAb

