

## MPP5 Rabbit pAb

CatalogNo: YN5822

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse

#### Applications

- WB

#### MW

- 74kD (Calculated)

#### Isotype

- IgG

### Recommended Dilution Ratios

WB 1:500-2000

### Storage

#### Storage\*

-15°C to -25°C/1 year(Do not lower than -25°C)

#### Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Basic Information

#### Clonality

Polyclonal

### Immunogen Information

#### Immunogen

Synthesized peptide derived from human MPP5

#### Specificity

This antibody detects endogenous levels of MPP5 at Human, Mouse

### Target Information

#### Gene name

MPP5

<b>Protein Name</b>	MAGUK p55 subfamily member 5		
	<b>Organism</b>	<b>Gene ID</b>	<b>UniProt ID</b>
	Human	<a href="#">64398</a> ;	<a href="#">Q8N3R9</a> ;
	Mouse	<a href="#">56217</a> ;	<a href="#">Q9JLB2</a> ;
<b>Cellular Localization</b>	<p>Golgi apparatus . Cell membrane; Peripheral membrane protein. Endomembrane system; Peripheral membrane protein. Cell junction, tight junction . Cell junction, adherens junction . Cell projection, axon . Perikaryon . Apical cell membrane . Localized to the tight junctions of epithelial cells (By similarity). Localized to the Golgi apparatus in T lymphocytes (PubMed:21479189). Localized to a subset of intracellular vesicles (By similarity). Localized to the Purkinje cell body and axon (By similarity). Localized to intercellular junctions in vascular endothelial cells (PubMed:27466317). Localized to Schmidt-Lanterman incisures, the adaxonal domain, and the inner part of paranodal loops in myelinating Schwann cells of the sciatic nerve (By similarity). Localized to apical membrane domains of the outer limiting membrane (OLM) junctions in the retina (By similarity). Colocalizes with CRB1 at the OLM, apical to the adherens junction (PubMed:15914641). Colocalizes with MPP1 in the retina at the OLM (PubMed:17584769). Colocalizes with MPP3 to the subapical region of adherens junctions in the retina OLM (PubMed:16519681). . ; Endoplasmic reticulum-Golgi intermediate compartment . Golgi apparatus . (Microbial infection) Following infection by human SARS coronavirus, partially localized at the site of viral replication; the endoplasmic reticulum-Golgi intermediate compartment, reducing its levels at cell-cell contacts which results in delayed formation of tight junctions and affects establishment of cell polarity. .</p>		
<b>Tissue specificity</b>	<p>Expressed at the outer limiting membrane in the retina (at protein level) (PubMed:15914641, PubMed:15558731, PubMed:16519681, PubMed:17584769). Expressed in T lymphocytes (at protein level) (PubMed:21479189). Expressed in the kidney (at protein level) (PubMed:17584769).</p>		
<b>Function</b>	<p>Plays a role in tight junction biogenesis and in the establishment of cell polarity in epithelial cells . Also involved in adherens junction biogenesis by ensuring correct localization of the exocyst complex protein EXOC4/SEC8 which allows trafficking of adherens junction structural component CDH1 to the cell surface (By similarity). Plays a role through its interaction with CDH5 in vascular lumen formation and endothelial membrane polarity . Required during embryonic and postnatal retinal development (By similarity). Required for the maintenance of cerebellar progenitor cells in an undifferentiated proliferative state, preventing premature differentiation, and is required for cerebellar histogenesis, fissure formation, cerebellar layer organization and cortical development (By similarity). Plays a role in neuronal progenitor cell survival, potentially via promotion of mTOR signaling (By similarity). Plays a role in the radial and longitudinal extension of the myelin sheath in Schwann cells (By similarity). May modulate SC6A1/GAT1-mediated GABA uptake by stabilizing the transporter (By similarity). Plays a role in the T-cell receptor-mediated activation of NF-kappa-B . Required for localization of EZR to the apical membrane of parietal cells and may play a role in the dynamic remodeling of the apical cytoskeleton (By similarity). Required for the normal polarized localization of the vesicular marker STX4 (By similarity). Required for the correct trafficking of the myelin proteins PMP22 and MAG (By similarity). Involved in promoting phosphorylation and cytoplasmic retention of transcriptional coactivators YAP1 and WWTR1/TAZ which leads to suppression of TGFB1-dependent transcription of target genes such as CCN2/CTGF, SERPINE1/PAI1, SNAI1/SNAI1 and SMAD7 (By similarity). . ; (Microbial infection) Acts as an interaction partner for human coronaviruses SARS-CoV and, probably, SARS-CoV-2 envelope protein E which results in delayed formation of tight junctions and dysregulation of cell polarity.</p>		

## | Validation Data

## | Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**MPP5 Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)