

## ST3GAL4 Rabbit pAb

CatalogNo: YN6488

## Key Features

Host SpeciesReactivityAp• Rabbit• Human,Mouse,Rat• VMWIsotype• 37kD (Calculated)• IgG

Applications
• WB

### Recommended Dilution Ratios

WB 1:500-2000

#### **Storage**

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### **Basic Information**

Clonality Polyclonal

#### Immunogen Information

Immunogen	Synthesized peptide derived from human ST3GAL4
Specificity	This antibody detects endogenous levels of ST3GAL4 at Human, Mouse,Rat

#### **Target Information**

Gene name ST3GAL4 CGS23 NANTA3 SIAT4C STZ

#### **Protein Name**

CMP-N-acetylneuraminate-beta-galactosamide-alpha-2,3-sialyltransferase 4 (Alpha 2,3-ST 4) (Beta-galactoside alpha-2,3-sialyltransferase 4) (Alpha 2,3-sialyltransferase IV) (Gal-NAc6S) (Gal-beta-1,4-GalNAc-alpha-2,3-sialyltransferase) (SAT-3) (ST-4) (ST3Gal IV) (ST3GalIV) (ST3GalA.2) (STZ) (Sialyltransferase 4C) (SIAT4-C)

Organism	Gene ID	UniProt ID
Human	<u>6484;</u>	<u>Q11206;</u>
Mouse	<u>20443;</u>	<u>Q91Y74;</u>
Rat	<u>363040;</u>	<u>P61131;</u>

# CellularGolgi apparatus, Golgi stack membrane; Single-pass type II membrane protein. Secreted.LocalizationMembrane-bound form in trans cisternae of Golgi. Secreted into the body fluid.

#### **Tissue specificity** Highly expressed in adult placenta, heart and kidney.

Function A beta-galactoside alpha2-3 sialyltransferase involved in terminal sialylation of glycoproteins and glycolipids. Catalyzes the transfer of sialic acid (N-acetyl-neuraminic acid; Neu5Ac) from the nucleotide sugar donor CMP-Neu5Ac onto acceptor Galbeta-(1->3)-GalNAc- and Galbeta-(1->4)-GlcNAc-terminated glycoconjugates through an alpha2-3 linkage . Plays a major role in hemostasis. Responsible for sialylation of plasma VWF/von Willebrand factor, preventing its recognition by asialoglycoprotein receptors (ASGPR) and subsequent clearance. Regulates ASGPR-mediated clearance of platelets (By similarity). Participates in the biosynthesis of the sialyl Lewis X epitopes, both on O- and N-glycans, which are recognized by SELE/E-selectin, SELP/P-selectin and SELL/L-selectin. Essential for selectin-mediated rolling and adhesion of leukocytes during extravasation. Contributes to adhesion and transendothelial migration of neutrophils likely through terminal sialylation of CXCR2 (By similarity). In glycosphingolipid biosynthesis, sialylates GM1 and GA1 gangliosides to form GD1a and GM1b, respectively. Metabolizes brain c-series ganglioside GT1c forming GQ1c (By similarity). Synthesizes ganglioside LM1 (IV3Neu5Ac-nLc4Cer), a major structural component of peripheral nerve myelin .

#### Validation Data

#### **Contact information**

- Orders: order.cn@immunoway.com
- Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

- Website: http://www.immunoway.com.cn
- Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: ST3GAL4 Rabbit pAb