

## Intersectin 1 Rabbit pAb

CatalogNo: YN6892

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB

#### MW

- 189kD (Calculated)

#### Isotype

- IgG

### Recommended Dilution Ratios

WB 1:500-2000

### Storage

**Storage\*** -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** Synthesized peptide derived from human Intersectin 1

**Specificity** This antibody detects endogenous levels of Intersectin 1 at Human, Mouse, Rat

### Target Information

**Gene name** ITSN1 ITSN SH3D1A

<b>Protein Name</b>	Intersectin-1 (SH3 domain-containing protein 1A) (SH3P17)		
	<b>Organism</b>	<b>Gene ID</b>	<b>UniProt ID</b>
	Human	<a href="#">6453</a> ;	<a href="#">Q15811</a> ;
	Mouse	<a href="#">16443</a> ;	<a href="#">Q9Z0R4</a> ;
	Rat	<a href="#">29491</a> ;	<a href="#">Q9WVE9</a> ;
<b>Cellular Localization</b>	Endomembrane system . Cell junction, synapse, synaptosome . Cell projection, lamellipodium . Cell membrane . Membrane, clathrin-coated pit . Recycling endosome . Endosome . Cytoplasmic vesicle . Colocalizes with SGIP1 at the plasma membrane in structures corresponding most probably to clathrin-coated pits (PubMed:20946875). Colocalizes with RAB13 on cytoplasmic vesicles that are most likely recycling endosomes (PubMed:29030480). .; [Isoform 2]: Cytoplasm . Endomembrane system . Nucleus envelope . Shuttles between the cytoplasm and nucleus in an XPO1/CRM1-dependent manner. .; [Isoform 5]: Endomembrane system .		
<b>Tissue specificity</b>	Isoform 1 is expressed almost exclusively in the brain. Isoform 2 is detected in brain, spleen, lung, liver, heart, skeletal muscle and kidney. Isoform 5 is primarily expressed in brain, spleen, lung and kidney (at protein level) (PubMed:21712076). Isoform 1 and isoform 2 are detected in brain (PubMed:10482960). Isoform 2 is ubiquitous in adult and fetal tissues with high expression in skeletal muscle, heart, spleen, ovary, testis and all fetal tissues tested and low expression in thymus, blood, lung, liver and pancreas. Isoform 1 is expressed almost exclusively in the brain, in all brain regions. Not expressed in the spinal cord (PubMed:9799604, PubMed:21712076, PubMed:11690630).		
<b>Function</b>	Adapter protein that provides a link between the endocytic membrane traffic and the actin assembly machinery . Acts as guanine nucleotide exchange factor (GEF) for CDC42, and thereby stimulates actin nucleation mediated by WASL and the ARP2/3 complex . Plays a role in the assembly and maturation of clathrin-coated vesicles (By similarity). Recruits FCHSD2 to clathrin-coated pits . Involved in endocytosis of activated EGFR, and probably also other growth factor receptors (By similarity). Involved in endocytosis of integrin beta-1 (ITGB1) and transferrin receptor (TFR); internalization of ITGB1 as DAB2-dependent cargo but not TFR may involve association with DAB2 . Promotes ubiquitination and subsequent degradation of EGFR, and thereby contributes to the down-regulation of EGFR-dependent signaling pathways. In chromaffin cells, required for normal exocytosis of catecholamines. Required for rapid replenishment of release-ready synaptic vesicles at presynaptic active zones (By similarity). Inhibits ARHGAP31 activity toward RAC1 .; [Isoform 1]: Plays a role in synaptic vesicle endocytosis in brain neurons.		

| Validation Data

| Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**Intersectin 1 Rabbit  
pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)