

IκB-β (Phospho Ser23) Rabbit pAb

CatalogNo: YP0153 **Orthogonal Validated** 

| Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 37kD (Observed)

Isotype

- IgG

| Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

| Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

IF 1:200-1:1000

ELISA 1:40000

Not yet tested in other applications.

| Basic Information

Clonality Polyclonal

| Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human IκappaB-beta around the phosphorylation site of Ser23. AA range:8-57

Specificity

Phospho-I κ B- β (S23) Polyclonal Antibody detects endogenous levels of I κ B- β protein only when phosphorylated at S23. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):LGSLG

Target Information

Gene name NFKB1B

Protein Name NF-kappa-B inhibitor beta


Organism	Gene ID	UniProt ID
Human	4793 ;	Q15653 ;
Mouse	18036 ;	Q60778 ;
Rat		Q9JIA3 ;

Cellular Localization Cytoplasm . Nucleus .


Tissue specificity Expressed in all tissues examined.

Function Function:Inhibits NF-kappa-B by complexing with and trapping it in the cytoplasm. However , the unphosphorylated form resynthesized after cell stimulation is able to bind NF-kappa-B allowing its transport to the nucleus and protecting it to further IKBA-dependent inactivation. Association with inhibitor kappa B-interacting NKIRAS1 and NKIRAS2 prevent its phosphorylation rendering it more resistant to degradation , explaining its slower degradation. ,PTM:Phosphorylated; followed by degradation. Interaction with NKIRAS1 and NKIRAS2 probably prevents phosphorylation. ,similarity:Belongs to the NF-kappa-B inhibitor family. ,similarity:Contains 6 ANK repeats. ,subunit:Interacts with THRB (via ligand-binding domain) . Interacts with RELA and REL. Interacts with COMMD1 and inhibitor kappa B-interacting Ras-like NKIRAS1 and NKIRAS2. ,tissue specificity:Expressed in all tissues examined. ,


Validation Data




Western Blot analysis of various cells using Phospho-IκB-β (S23) Polyclonal Antibody




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using IκappaB-beta (Phospho-Ser23) Antibody



Immunofluorescence analysis of HeLa cells treated with TNF-a 20nM 15', using IκappaB-beta (Phospho-Ser23) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma, using IκappaB-beta (Phospho-Ser23) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HeLa cells treated with TNF- α 20ng/ml 5', using I κ B- β (Phospho-Ser23) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
I κ B- β (Phospho Ser23) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)