

MEK-6 (Phospho Ser207) Rabbit pAb

CatalogNo: YP0173 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 37kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:40000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized phospho-peptide around the phosphorylation site of human MEK-6 (phospho Ser207)

Specificity Phospho-MEK-6 (S207) Polyclonal Antibody detects endogenous levels of MEK-6 protein only when phosphorylated at S207. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):VDsVA

Target Information

Gene name MAP2K6

Protein Name Dual specificity mitogen-activated protein kinase kinase 6


Organism	Gene ID	UniProt ID
Human	5608 ;	P52564 ;
Mouse	26399 ;	P70236 ;

Cellular Localization Nucleus . Cytoplasm . Cytoplasm , cytoskeleton . Binds to microtubules.

Tissue specificity Isoform 2 is only expressed in skeletal muscle. Isoform 1 is expressed in skeletal muscle , heart , and in lesser extent in liver or pancreas.

Function Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,enzyme regulation:Probably activated by dual phosphorylation on Ser-207 and Thr-211. ,Function:Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in MAP kinase p38 exclusively. ,induction:Strongly activated by UV , anisomycin , and osmotic shock but not by phorbol esters , NGF or EGF. ,PTM:Acetylation of Ser-207 and Thr-211 by Yersinia yopJ prevents phosphorylation and activation , thus blocking the MAPK signaling pathway. ,PTM:Weakly autophosphorylated. ,similarity:Belongs to the protein kinase superfamily. ,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase subfamily. ,similarity:Contains 1 protein kinase domain. ,subunit:Interacts with Yersinia yopJ. ,tissue specificity:Isoform 2 is only expressed in skeletal muscle. Isoform 1 , on the other hand , is found in skeletal muscle , heart , and in lesser extent in liver or pancreas. ,

Validation Data



Immunohistochemical analysis of paraffin-embedded human small intestinal carcinoma tissue. 1,primary Antibody was diluted at 1:200 (4°C overnight). 2, Sodium citrate pH 6.0 was used for antigen retrieval (>98°C,20min). 3,Secondary antibody was diluted at 1:200

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
MEK-6 (Phospho Ser207) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)