

GSK3 β (Phospho Ser9) Rabbit pAb

CatalogNo: YP0285 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 47kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:10000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized phospho-peptide around the phosphorylation site of human GSK3 β (phospho Ser9)

Specificity

Phospho-GSK3 β (S9) Polyclonal Antibody detects endogenous levels of GSK3 β protein only when phosphorylated at S9. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): TTsFA

Target Information

Gene name GSK3B

Protein Name Glycogen synthase kinase-3 beta

Organism	Gene ID	UniProt ID
Human	2932 ;	P49841 ;
Mouse	56637 ;	Q9WV60 ;
Rat	84027 ;	P18266 ;

Cellular Localization

Cytoplasm . Nucleus . Cell membrane . The phosphorylated form shows localization to cytoplasm and cell membrane (PubMed:20937854) . The MEMO1-RHOA-DIAPH1 signaling pathway controls localization of the phosphorylated form to the cell membrane (PubMed:20937854) . .


Tissue specificity

Expressed in testis , thymus , prostate and ovary and weakly expressed in lung , brain and kidney. Colocalizes with EIF2AK2/PKR and TAU in the Alzheimer disease (AD) brain.


Function

Catalytic activity:ATP + [tau protein] = ADP + [tau protein] phosphate. ,enzyme regulation:Inhibited when phosphorylated by AKT1. ,Function:Participates in the Wnt signaling pathway. Implicated in the hormonal control of several regulatory proteins including glycogen synthase , MYB and the transcription factor JUN. Phosphorylates JUN at sites proximal to its DNA-binding domain , thereby reducing its affinity for DNA. Phosphorylates MUC1 in breast cancer cells , and decreases the interaction of MUC1 with CTNNB1/beta-catenin. ,PTM:Phosphorylated by AKT1 and ILK1. ,similarity:Belongs to the protein kinase superfamily. ,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. GSK-3 subfamily. ,similarity:Contains 1 protein kinase domain. ,subunit:Monomer (By similarity) . Interacts with CABYR , MUC1 , NIN and PRUNE. ,tissue specificity:Expressed in testis , thymus , prostate and ovary and weakly expressed in lung , brain and kidney. ,

Validation Data



Western Blot analysis of RAT-kidney cells using Phospho-GSK3β (S9) Polyclonal Antibody diluted at 1:500



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
GSK3β (Phospho Ser9) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)