

www.immunoway.com.cn

c-Fms (Phospho Tyr809) Rabbit pAb

CatalogNo: YP0308 Orthogonal Validated 💽

Key Features

| Host Species Rabbit | Reactivity • Human,Mouse,Rat | ApplicationsWB,ELISA |
|---|---------------------------------|---|
| MW • 130-170kD (Observed) | Isotype • IgG | |

Recommended Dilution Ratios

WB 1:500-1:2000 ELISA 1:5000 Not yet tested in other applications.

Storage

| Storage* | -15°C to -25°C/1 year(Do not lower than -25°C) |
|-------------|---|
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |

Basic Information

Clonality Polyclonal

Immunogen Information


- **Immunogen** The antiserum was produced against synthesized peptide derived from human M-CSF Receptor around the phosphorylation site of Tyr809. AA range:781-830
- **Specificity** Phospho-c-Fms (Y809) Polyclonal Antibody detects endogenous levels of c-Fms protein only when phosphorylated at Y809. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):SNyVV

Target Information

| Gene name | CSF1R | | |
|--------------------------|---|------------------------|----------------------|
| Protein Name | Macrophage colony-stimulating factor 1 receptor | | |
| | Organism | Gene ID | UniProt ID |
| | Human | <u>1436;</u> | <u>P07333;</u> |
| | Mouse | <u>12978;</u> | <u>P09581;</u> |
| | Rat | | <u>Q00495;</u> |
| Cellular Localization | Cell membrane; Single-pass type I membrane protein. | | |
| Tissue specificity | Expressed in bone marrow and in | n differentiated blood | l mononuclear cells. |
| Function | Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,Function:Protein tyrosine-kinase transmembrane receptor for CSF1 and IL34.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. CSF-1/PDGF receptor subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 5 Ig-like C2-type (immunoglobulin-like) | | |


bone marrow and in differentiated blood mononuclear cells.,

Validation Data



Western blot analysis of 293 using p-c-Fms (Y809) antibody.

domains., subunit: Interacts with INPPL1/SHIP2 and THOC5., tissue specificity: Expressed in



Western blot analysis of lysates from 293 cells treated with LPS 100ng/ml 30', using M-CSF Receptor (Phospho-Tyr809) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

| Orders: | order.cn@immunoway.com |
|------------|--|
| Support: | support.cn@immunoway.com |
| Telephone: | 400-8787-807(China) |
| Website: | http://www.immunoway.com.cn |
| Address: | 2200 Ringwood Ave San Jose, CA 95131 USA |
| | |



Please scan the QR code to access additional product information: **c-Fms (Phospho Tyr809) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents