

Survivin (Phospho Thr34) Rabbit pAb

CatalogNo: YP0319 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA, IHC

MW

- 16kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

IHC 1:50-300

ELISA 1:2000-20000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human Survivin around the phosphorylation site of Thr34. AA range:11-60

Specificity Phospho-Survivin (T34) Polyclonal Antibody detects endogenous levels of Survivin protein only when phosphorylated at T34. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):ACTPE

Target Information

Gene name BIRC5

Protein Name Baculoviral IAP repeat-containing protein 5

| Organism | Gene ID | UniProt ID |
|----------|-------------------------|--------------------------|
| Human | 332 ; | O15392 ; |
| Mouse | 11799 ; | O70201 ; |
| Rat | 64041 ; | Q9JHY7 ; |

Cellular Localization

Cytoplasm . Nucleus . Chromosome . Chromosome , centromere . Cytoplasm , cytoskeleton , spindle . Chromosome , centromere , kinetochore . Midbody . Localizes at the centromeres from prophase to metaphase , at the spindle midzone during anaphase and at the midbody during telophase and cytokinesis. Accumulates in the nucleus upon treatment with leptomycin B (LMB) , a XPO1/CRM1 nuclear export inhibitor (By similarity) . Localizes on chromosome arms and inner centromeres from prophase through metaphase. Localizes to kinetochores in metaphase , distributes to the midzone microtubules in anaphase and at telophase , localizes exclusively to the midbody (PubMed:11084331) . Colocalizes with AURKB at mitotic chromosomes (PubMed:14610074) . Acetylation at Lys-129 directs its localization to the nucleus by enhancing homodimerization and thereby inhibiting XPO1/CRM1-mediated nuclear export (PubMed:20826784) . .


Tissue specificity

Expressed only in fetal kidney and liver , and to lesser extent , lung and brain (PubMed:10626797) . Abundantly expressed in adenocarcinoma (lung , pancreas , colon , breast , and prostate) and in high-grade lymphomas (PubMed:14741722 , PubMed:16329164) . Also expressed in various renal cell carcinoma cell lines (PubMed:10626797) . Expressed in cochlea including the organ of Corti , the lateral wall , the interdental cells of the Limbus as well as in Schwann cells and cells of the cochlear nerve and the spiral ganglions (at protein level) . Not expressed in cells of the inner and outer sulcus or the Reissner's membrane (at protein level) (PubMed:21364656 , PubMed:20627126) .


Function

Domain:The BIR repeat is necessary and sufficient for HBXIP binding. ,Function:May play a role in neoplasia. May counteract a default induction of apoptosis in G2/M phase. Interacts with tubulin. Inhibitor of caspase-3 and caspase-7. Component of the chromosomal passenger complex (CPC) , a complex that acts as a key regulator of mitosis. The CPC complex has essential functions at the centromere in ensuring correct chromosome alignment and segregation and is required for chromatin-induced microtubule stabilization and spindle assembly. Isoforms 2 and 3 do not appear to play vital roles in mitosis. Isoform 3 shows a marked reduction in its anti-apoptotic effects when compared with the displayed wild-type isoform. ,similarity:Belongs to the IAP family. ,similarity:Contains 1 BIR repeat. ,subcellular location:Localizes on chromosome arms and inner centromeres from prophase through metaphase and then transferring to the spindle midzone and midbody from anaphase through cytokinesis. Colocalizes with AURKB at mitotic chromosomes. ,subunit:Homodimer. When phosphorylated , interacts with HBXIP; the resulting complex binds pro-caspase-9 , as well as active caspase-9 , but much less efficiently. Component of the CPC at least composed of BIRC5/survivin , CDCA8/borealin , INCENP and AURKB/Aurora-B. Interacts with EVI5. ,tissue specificity:Expressed only in fetal kidney and liver , and to lesser extent , lung and brain. Abundantly expressed in adenocarcinoma (lung , pancreas , colon , breast , and prostate) and in high-grade lymphomas. Also expressed in various renal cell carcinoma cell lines. ,

Validation Data



Western blot analysis of lysates from HT29 cells treated with EGF 100ug/ml 30', using Survivin (Phospho-Thr34) Antibody. The lane on the right is blocked with the phospho peptide.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Survivin (Phospho Thr34) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)